



# First Aid Made Easy

A comprehensive first aid  
manual and reference guide.

|                     |     |
|---------------------|-----|
| AED Usage           | 58  |
| Amputation          | 32  |
| Anaphylaxis         | 18  |
| Angina              | 24  |
| Asthma              | 19  |
| Bleeding            | 30  |
| Burns               | 34  |
| Chest Injury        | 21  |
| Choking             | 16  |
| CPR                 | 6-8 |
| Croup               | 19  |
| Crush Injury        | 32  |
| Diabetes            | 46  |
| Drowning            | 20  |
| Epilepsy            | 48  |
| Eye Injury          | 31  |
| Fainting            | 27  |
| Febrile Convulsions | 49  |
| Fitting             | 48  |
| Fractures           | 37  |
| Frostbite           | 44  |
| Head Injuries       | 12  |
| Heart Attack        | 24  |
| Heat Exhaustion     | 44  |
| Heat Stroke         | 45  |
| Hyperventilation    | 20  |
| Hypothermia         | 43  |
| Internal Bleeding   | 32  |
| Poisoning           | 33  |
| Recovery Position   | 11  |
| Resuscitation       | 6-8 |
| Shock               | 26  |
| Spinal Injury       | 39  |
| Sprains and Strains | 37  |
| Stroke              | 13  |
| Unconsciousness     | 9   |



## Contents

|                                      |       |
|--------------------------------------|-------|
| <b>First Aid</b>                     |       |
| Introduction                         | 3     |
| The Aims of First Aid                | 4     |
| Priorities of Treatment              | 4     |
| <b>Resuscitation</b>                 |       |
| Emergency Action Plan                | 5     |
| Chain of Survival                    | 6     |
| Cardio Pulmonary Resuscitation (CPR) | 6-8   |
| Child and Baby CPR                   | 8     |
| Chest Compression only CPR           | 8     |
| <b>Unconsciousness</b>               |       |
| Definition and Causes                | 9     |
| Levels of Response                   | 9     |
| Primary and Secondary Survey         | 10    |
| Recovery Position                    | 11    |
| Head Injuries                        | 12-13 |
| Stroke                               | 13    |
| Hypoxia (low oxygen)                 | 14    |
| <b>Airway and Breathing Problems</b> |       |
| The Respiratory System               | 15    |
| Choking                              | 16-17 |
| Anaphylaxis                          | 18    |
| Asthma and Croup                     | 19    |
| Hyperventilation                     | 20    |
| Drowning                             | 20    |
| Chest Injuries                       | 21    |
| <b>Circulation Problems</b>          |       |
| The Circulatory System               | 22-23 |
| Angina                               | 24-25 |
| Heart Attack                         | 24-25 |
| Left Ventricular Failure             | 25    |
| Shock: - Hypovolaemic                | 26    |
| - Cardiogenic                        | 27    |
| - Anaphylactic                       | 27    |
| Fainting                             | 27    |
| <b>Wounds and Bleeding</b>           |       |
| Hygiene                              | 28    |
| Types of Wound                       | 28    |
| Blood Loss                           | 29    |
| Treatment of Bleeding                | 30    |
| Embedded Objects                     | 31    |
| Nose Bleeds                          | 31    |
| Eye Injuries                         | 31    |
| Amputation                           | 32    |
| Crush Injury                         | 32    |
| Internal Bleeding                    | 32    |

|  |       |
|--|-------|
| <b>Poisons, Burns and Scalds</b>             |       |
| Poisons                                      | 33    |
| Estimating Severity of a Burn                | 34    |
| Causes of Burns and Treatment                | 35    |
| <b>Injuries to Bones, Muscles and Joints</b> |       |
| The Skeletal System                          | 36    |
| Causes of Injury                             | 37    |
| Types of Fracture                            | 37    |
| Dislocations                                 | 37    |
| Sprains and Strains                          | 37    |
| Sprains and Strains – Treatment              | 39    |
| Fractures – Signs and Symptoms               | 38    |
| Fractures – Treatment                        | 38    |
| Spinal Injuries                              | 39-41 |
| <b>Effects of Heat and Cold</b>              |       |
| Body Temperature                             | 42    |
| Hypothermia                                  | 43    |
| Frostbite                                    | 44    |
| Heat Exhaustion                              | 44    |
| Heat Stroke                                  | 45    |
| Taking a Temperature                         | 45    |
| <b>Other Serious Conditions</b>              |       |
| Diabetes                                     | 46-47 |
| Epilepsy                                     | 48-49 |
| Seizures                                     | 48-49 |
| Febrile Convulsions                          | 49    |
| The Digestive System                         | 50    |
| <b>Health and Safety</b>                     |       |
| Employer's Responsibilities                  | 51    |
| First Aid Kits                               | 51    |
| First Aid needs assessment                   | 52    |
| Typed of <b>First Aider</b>                  | 53    |
| Reporting of Incidents (RIDDOR)              | 53    |
| Accident Book                                | 54    |
| Patient Report Form                          | 55    |
| <b>Appendix</b>                              |       |
| Resuscitation Child                          | 56    |
| Resuscitation Baby                           | 57    |
| Resuscitation with an AED                    | 58-60 |
| <b>Glossary</b>                              |       |
| Notes  | 61-64 |
| Glossary                                     | 65    |
| <b>First Aid Quiz</b>                        |       |
| First Aid Quiz                               | 66    |

## Introduction

*This manual has been designed by an experienced paramedic instructor to guide you through your first aid course, and to provide you with a reference for future years.*

*Most people will find the information in this book useful and informative, but it cannot replace 'hands on' training in the vital skills of dealing with an emergency situation.*

*Effective emergency treatment before professional help arrives can go a long way to reducing the effects of illness and injury, and indeed save someone's life.*

*Taking part in a first aid course and using this manual may be the most important decision you make in your life...*

### Edition 7.1

#### IMPORTANT

This manual is designed as a learning guide to a full first aid course, it cannot replace 'hands on' training in the vital skills of dealing with an emergency situation.

If you suspect illness or injury, you should always seek professional medical advice.

#### DISCLAIMER

Whilst every effort has been made to ensure the accuracy of the information contained within this manual, the author does not accept any liability for any inaccuracies or for any subsequent mistreatment of any person, however caused.

© 2008. Quallsafe Limited. All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the copyright owner.  
Tel: 0845 644 3305 www.quallsafe.com

## The aims of first aid

### Preserve Life

Not only the casualty's life, but your own as well. Far too often only one person's life is in danger when the emergency services are called, but by the time they arrive there are more. If you put your life in danger, you can end up fighting for your OWN life instead of the casualty's.

### Prevent the situation from Worsening

The skilled first aider must take action to prevent the whole situation from becoming worse (*e.g. removing dangers such as traffic or fumes*), as well as acting to prevent the casualty's condition from deteriorating.

### Promote Recovery

The actions of a first aider should, after preventing things from getting worse, help the casualty to recover from their illness or injury.


## Priorities of treatment

The first priority with any patient is to make sure the Airway is open and then to check they are Breathing normally (*A and B*). If the patient is breathing normally, this means that their heart must also be beating, so blood is being circulated around the body. As the A and B check is carried out first, we call it the '*primary survey*'.

Once you are happy that the casualty is Breathing normally and oxygen is being circulated around the body, the next priority is to deal with any major Bleeding, because you need to maintain enough blood to circulate the oxygen around. After these steps, the next priority is to deal with any broken Bones (*BBB*). The check for bleeding and then broken bones is called the '*secondary survey*'.

*All animal life needs a constant supply of oxygen to survive. If that oxygen is taken away for any reason, brain cells will start to die within 3 to 4 minutes.*


*The priorities of treatment are therefore aimed firstly at getting oxygen into the blood stream, ensuring that the blood is circulating around the body, and then preventing the loss of that blood. If this aim is achieved, then the majority of casualties will still be alive when the ambulance arrives.*



### Multiple casualties

The **BBB** rule can be used for multiple casualties, to decide who needs treatment first. A rough 'rule of thumb' is that the casualty who is the quietest needs treatment first, where as the one making the most noise (trying to get your attention) is the least serious!



It is important to have an action plan for emergencies. This flow chart guides you through the actions to be taken when dealing with a patient. All the topics, such as the recovery position and resuscitation are covered later in the book.



## The chain of survival

In order to maintain the oxygen supply to the body a person must be breathing, and their heart must be 'pumping'. If either of these two functions stop, the brain and other vital organs will quickly deteriorate, and brain cells will start to die within 3 to 4 minutes. Unless urgent action is taken to circulate oxygen around the body, this will inevitably result in death.

The most common cause of cardiac arrest in adults is 'ventricular fibrillation'. In these circumstances the best chance of restarting the heart is by using a 'defibrillator', which is carried on all emergency ambulances in the UK. For this reason, an emphasis is placed on summoning help and dialling 999 as soon as possible. Of course, the heart and brain must be kept oxygenated until the defibrillator arrives; so early Cardio Pulmonary Resuscitation (CPR) is vital if a casualty is to recover. These actions form the 'links' in the chain of survival (see diagram).



## Cardio Pulmonary Resuscitation (CPR) – Primary survey:

### **D** Danger – *make sure it's safe and find out what's happened*

- Check that it is safe for you to help the casualty. Do not put yourself at risk in any way.
- If possible remove any danger from the casualty, or if not, can you safely move the casualty from the danger?
- Find out what's happened – and make sure you are still safe.
- Check how many casualties there are. Can you cope?

### **R** Response – *are they conscious?*

- Gently shake the shoulders and ask loudly 'Are you alright?'
- If there is no response, shout for help immediately, but do not leave the casualty yet.

### **A** Airway – *open the airway*

- Carefully open the airway by using 'head tilt' and 'chin lift':
  - Place your hand on the forehead and gently tilt the head back.
  - With your fingertips under the point of the casualty's chin, lift the chin to open the airway (see diagram).

### **B** Breathing – *check for normal breathing*

Keeping the airway open, check to see if the breathing is normal. Take no more than 10 seconds to do this:

- Look at the chest and abdomen for movement.
- Listen for the sounds of breathing (*more than the occasional gasp*).
- Feel for air on your cheek or movement of the chest or abdomen.

If the casualty is breathing normally, carry out a secondary survey and place them in the recovery position (pages 11 and 12).



Gently shake the shoulders and shout.



Gently tip the head back and lift the chin to open the airway.

**WARNING:** In the first few minutes after cardiac arrest, a casualty may be barely breathing, or taking infrequent, noisy, gasps. Do not confuse this with normal breathing. If you are in doubt, start CPR.

**If the casualty is not breathing normally:**

Ask someone to **dial 999** for an ambulance or, if you are on your own, do this yourself; you may need to leave the casualty. Start chest compressions as follows:

- Place the heel of one hand in the centre of the casualty's chest, then place the heel of your other hand on top and interlock your fingers (*see diagram*).
- Position yourself vertically above the casualty's chest with your arms straight.
- Press down on the breastbone 4 to 5cm (*1 1/2 to 2 inches*) then release the pressure without losing contact between your hands and the chest (*chest compression*). Ensure that pressure is not applied over the casualty's ribs. Don't apply pressure over the upper abdomen or the bottom end of the breastbone.
- Compression and release should take an equal amount of time.
- Do 30 chest compressions at a rate of 100 per minute.
- Now combine chest compressions with rescue breaths (*below*).

*NOTE: Ideally the casualty needs to be on a firm flat surface to perform chest compressions (not a bed). One way to remove someone from a low bed is to unhook the bed sheets and use them to slide the casualty carefully to the floor. Get help if you can and be very careful not to injure yourself or the casualty. Do not move the casualty if you do not think it's safe to do so – remove the pillows and attempt CPR on the bed instead.*

**Combine chest compression with rescue breaths:**

- Open the airway again, using head tilt and chin lift.
- Nip the soft part of the casualty's nose closed. Allow the mouth to open, but maintain chin lift.
- Take a normal breath and seal your lips around the casualty's mouth.
- Blow steadily into the casualty's mouth, whilst watching for the chest to rise (*rescue breath*). Take about one second to make the chest rise.
- Keeping the airway open, remove your mouth. Take a breath of fresh air and watch for the casualty's chest to fall as air comes out.
- Re-seal your mouth and give another rescue breath (*two in total*).
- Return your hands without delay to the correct position on the breastbone and give another 30 chest compressions (*then 2 more rescue breaths*).
- **Continue repeating cycles of 30 chest compressions and 2 rescue breaths.**
- Only stop to recheck the casualty if they start breathing normally – otherwise don't interrupt resuscitation.

If your rescue breaths don't make the chest rise effectively, give another 30 chest compressions, then before your next attempt:

- Check the casualty's mouth and remove any visible obstruction.
- Recheck that there is adequate head tilt and chin lift.
- Do not attempt more than two breaths each time before returning to chest compressions.

*NOTE: If there is more than one rescuer, change over every two minutes to prevent fatigue. Ensure the minimum of delay as you change over.*

**Continue resuscitation until:**

- Qualified help arrives and takes over.
- The casualty starts breathing normally, or
- You become exhausted.



*Look, listen and feel for normal breathing.*



*Place the heel of one hand in the centre of the chest, then the other hand on top.*



*Arms straight and shoulders above your hands, depress the chest 4 to 5cm.*



*Nip the nose and seal your mouth around the casualty's mouth.*



*Slowly breathe just enough air to make the chest rise.*



For a child over 1 year, use one or two hands to compress the chest by about one third of its depth.



For a baby under 1 year, use two fingers to compress the chest by about one third of its depth.



Turn them onto their side and allow the vomit to run out.



Using a protective barrier during CPR.

## Resuscitation for children and babies



Recent studies have found that many children do not receive resuscitation because potential rescuers fear causing them harm. It is important to understand that it's far better to perform 'adult style' resuscitation on a child (*who is unresponsive and not breathing*) than to do nothing at all.

For ease of learning and retention, first aiders can use the adult sequence of resuscitation (*see previous pages*) on a child or baby who is unresponsive and not breathing. The following minor modifications to the adult sequence will, however, make it even more suitable for use in children:

- Give five initial rescue breaths before starting chest compressions (*then continue at the ratio of 30 compressions to 2 breaths*).
- If you are on your own, perform resuscitation for about 1 minute before going for help.
- Compress the chest by about one-third of its depth:
  - For a baby under 1 year, use two fingers.
  - For a child over 1 year, use one or two hands (*as needed*) to achieve an adequate depth of compression (*about one third of the depth*).

The full sequence of child or baby resuscitation is given in detail on pages 56 and 57.

## Chest compression only resuscitation



When an **adult** casualty suffers a cardiac arrest, it is likely that there is residual oxygen left in the blood stream.

If you are unable (*or unwilling*) to give rescue breaths, give 'chest compressions only' resuscitation, as this will circulate any residual oxygen in the blood stream, so it is better than no CPR at all.

- If chest compressions only are given, these should be continuous at a rate of 100 per minute.
- Stop to recheck the casualty only if they start breathing normally – otherwise do not interrupt resuscitation.
- If there is more than one rescuer, change over every two minutes to prevent fatigue. Ensure the minimum of delay as you change over.

## Vomiting



It is common for a patient who has stopped breathing to vomit whilst they are collapsed. This is a passive action in the unconscious person, so you may not hear or see it happening. You might not find out until you give a rescue breath (*as the air comes back out of the patient it makes gurgling noises*).

- If the patient has vomited, turn them onto their side, tip the head back and allow the vomit to run out.
- Clean the face of the patient then continue resuscitation, using a protective face barrier if possible.

## Hygiene during resuscitation:

- Wipe the lips clean.
- If possible use a protective barrier such as a 'face shield'. (*This is particularly important if the patient suffers from any serious infectious disease such as TB, Hepatitis or S.A.R.S.*).
- As a last resort some plastic with a hole in it, or a handkerchief, may help to prevent direct contact.
- If you are still in doubt about the safety of performing rescue breaths, give 'chest compression only' resuscitation (*see above*).
- Wear protective gloves if available and wash your hands afterwards.



## The main causes of unconsciousness

The causes of unconsciousness can be remembered by using 'FISH SHAPED.' Each of these causes are dealt with individually elsewhere in this manual.

- |                           |                       |
|---------------------------|-----------------------|
| <b>F</b> ainting          | <b>S</b> troke        |
| <b>I</b> mbalance of heat | <b>H</b> earth Attack |
| <b>S</b> hock             | <b>A</b> sphyxia      |
| <b>H</b> ead Injury       | <b>P</b> oisoning     |
|                           | <b>E</b> pilepsy      |
|                           | <b>D</b> iabetes      |

Unconsciousness can be defined as an interruption in the normal activity of the brain. Unlike sleep, unconsciousness can disable the body's natural reflexes such as coughing. Therefore if the unconscious patient is laying on their back the tongue may fall back blocking the airway, or they may even drown themselves if they vomit.

You should take immediate action to treat an unconscious casualty. This will involve protecting the airway, calling an ambulance and possibly treating the underlying cause of the condition.

## Levels of response

In order to accurately measure a casualty's conscious level, we can use a scale of consciousness called the 'AVPU' scale:

|                      |  |
|----------------------|--|
| <b>A</b> lert        | <b>The casualty is fully alert</b><br>They are responsive and fully orientated (a casualty in this category will usually know what month it is). |
|                      | <b>Confused</b><br>The casualty is not fully orientated but asks and answers your questions.   |
|                      | <b>Inappropriate Words</b><br>The patient is able to speak words, but cannot put them together into logical sentences.                           |
|                      | <b>Utters Sounds</b><br>The casualty is not able to speak words but makes noises, often in response to painful stimuli.                          |
| <b>V</b> oice        | <b>No Verbal Response</b><br>The casualty makes no noise.  |
|                      | <b>Localises Pain</b><br>The patient is able to localise where painful stimuli is being applied.   |
|                      | <b>Responds to (but does not localise) Pain</b><br>The patient responds to painful stimuli, but is unable to localise it.                        |
| <b>P</b> ain         | <b>Unresponsive</b><br>The casualty is unresponsive to pain and speech stimuli.  |
| <b>U</b> nresponsive |  |



# Unconsciousness

The Primary and Secondary Survey methods of checking a patient give us a systematic order in which to deal with the most urgent problems first, then move on to find other clues – helping with diagnosis and treatment.

## Primary survey

When you check for Danger, Response, Airway and Breathing this is called the 'Primary Survey.' This can be found in the 'resuscitation' section of this manual (see page 6).

The primary survey ensures that the patient is breathing, so it should be carried out first.


Once you are sure that the patient is breathing effectively, it is safe to move on and carry out a secondary survey:

## Secondary survey

If a casualty is unconscious you are concerned about the airway for any reason (e.g. vomiting), place them in the recovery position immediately (page 11).

The Secondary Survey should be done quickly and systematically, first checking for major bleeding and then broken bones.

- Bleeding**
- Do a quick head to toe check for bleeding.
  - Check the hidden area such as under the arch of the back.
  - Control any major bleeding that you find (page 30).
- Head and neck**
- Clues to injury could be bruising, swelling, deformity or bleeding.
  - Check the whole head and face.
  - Feel the back of the neck.
  - Has the patient had an accident that might have injured the neck? (page 39).
- Shoulders and chest**
- Place your hands on opposite shoulders and compare them.
  - Run your fingers down the collar bones checking for signs of a fracture (page 38).
  - Gently squeeze and rock the ribs.
- Abdomen and pelvis**
- Push the abdomen with the palm of your hand to check for abnormality or response to pain.
  - Gently check the pelvis for signs of a fracture.
  - Look for incontinence or bleeding.
- Legs and arms**
- Feel each leg for the signs of a fracture.
  - Feel each arm for the signs of a fracture.
  - Look for other clues (medic alert bracelets, needle marks etc).
- Pockets**
- Look for clues and make sure nothing will injure the patient as you roll them into the recovery position.
  - Have a witness if you remove items from pockets.
  - Be very careful if you suspect there could be sharp objects such as needles.
  - Loosen any tight clothing.
- Recovery**
- Place the patient in the recovery position (page 11).
  - If you suspect neck injury, get someone to help you keep the head in line with the body as you turn the patient (see page 41 for how to do this).
  - Be careful not to cause further damage to any suspected injuries.



Remember the priorities of treatment? (page 4)

## Mechanics of injury

Before you move a patient, it is important to consider the 'mechanics of injury'.

This involves trying to work out what happened and what injuries this could have caused the patient.

- If you suspect neck injury, get someone to help you keep the head in line with the body at all times (see page 41 for how to do this).
- If you have to use the recovery position, try not to move any suspected injuries.

## The recovery position



When a person is unconscious and lying on their back, the airway can become compromised by the tongue touching the back of the throat, or vomit if the patient is sick. Placing the casualty in the recovery position protects the airway from both of these dangers – the tongue will not fall backwards and vomit will run out of the mouth.

1



- Remove the casualty's glasses and straighten both legs.
- Move the arm nearest you outwards, elbow bent with palm uppermost.

2



- Bring the far arm across the chest, and hold the back of that hand against the cheek.

3



- With your other hand, grasp the far leg just above the knee, and pull it up, keeping the foot on the ground.
- Keeping the casualty's hand pressed against their cheek, pull on the leg to roll them towards you, onto their side.

4



- Adjust the upper leg so that the hip and knee are bent at right angles and tilt the head back to keep the airway open.
- Call 999 for an ambulance.
- Check breathing regularly. If breathing stops, turn the casualty onto their back again and perform CPR.

**NEVER** place anything in an unconscious casualty's mouth.

**NEVER** move a casualty without checking them first.

**NEVER** place a pillow under the head whilst the casualty is on their back.

**NEVER** move the casualty unnecessarily.

## Hypoxia

The medical term 'hypoxia' means 'low oxygen in the blood'.

A low level of oxygen in the blood is potentially fatal, so it is very important that the First Aider recognises the signs and symptoms of this condition and takes immediate action to treat the casualty.

The causes of hypoxia can be separated into 5 areas:

### Possible signs and symptoms ?

- *Pale clammy skin (for dark skinned casualties look at the colour of skin inside the lips).*
- *Blue tinges to the skin and lips (cyanosis).*
- *Increase in pulse rate.*
- *Weakening of the pulse.*
- *Nausea or vomiting.*
- *Increased breathing rate (caused by oxygen deficiency).*
- *Lowered breathing rate (look for control centre causes).*
- *Distressed breathing or gasping.*
- *Confusion or dizziness.*
- *Lowering levels of consciousness.*
- *Clues from the cause of the hypoxia (i.e. bleeding, injury, chest pain etc).*



*Pale clammy skin and cyanosis.*

### Treatment of hypoxia +

- *Maintain **Airway** and **Breathing** (pages 6 to 8).*
- *Remove or treat the **cause** of the hypoxia.*
- *Do not allow the patient to eat, drink or smoke.*

### External causes:

Not enough oxygen in the surrounding air, such as:

- Suffocation by gas or smoke.
- Suffocation by sand, earth or a pillow etc.
- Drowning.
- High altitude.

### Airway causes:

Blockage, swelling or narrowing. Caused by:

- The tongue.
- Strangulation.
- Vomit.
- Hanging.
- Choking.
- Anaphylaxis.
- Burns.

### Breathing causes:

Inability of the lungs to function properly. Caused by:

- Crushing of the chest.
- Poisoning.
- Collapsed lung.
- Asthma.
- Chest injury.
- Disease or illness.

### Circulation causes:

Inability of the blood to take up oxygen, a fall in blood pressure, or failure to circulate the blood around the body. Caused by:

- Heart attack.
- Severe bleeding.
- Cardiac arrest.
- Poisoning.
- Angina.
- Anaemia.

### Control centre causes:

Failure of the respiratory control centre in the brain or the nerves connecting it to the lungs. Caused by:

- Stroke.
- Poisoning.
- Head injury.
- Spinal injury.
- Drug overdose.
- Electric shock.

## The body's response to hypoxia



If the body detects low levels of oxygen in the blood ADRENALINE is released. The effect of adrenaline on the body is to:

- Increase the heart rate.
- Increase the strength of the heart beat (*and blood pressure*).
- Divert blood away from the skin, intestines and stomach.
- Divert blood towards the heart, lungs and brain.
- Dilate the air passages in the lungs (*bronchioles*).

The effect of adrenaline being released in the body produces dramatic signs and symptoms that the first aider must be able to recognise.

Can you tell which of the signs and symptoms are caused by adrenaline?

## The respiratory system



Oxygen passes from the alveoli into the blood, whilst carbon dioxide passes in the opposite direction to be breathed out.

Air is drawn in through the mouth and nose, where it is warmed, filtered and moistened.

Air then travels through the throat and past the epiglottis (the protective flap of skin that folds down to protect the airway when we swallow).

Air now enters the larynx (more commonly known as the voice box or 'Adam's apple'). It passes between the vocal cords in the larynx and down into the trachea.

The trachea is protected by rings of cartilage that surround it to prevent kinking.

The trachea divides into two 'bronchi' that supply air to each lung.

The bronchi then divide into smaller air passages called 'bronchioles'.

At the end of the bronchioles are microscopic air sacks called 'alveoli'.

The walls of the alveoli are only one cell in thickness, so oxygen can pass through into the blood, which is carried in capillaries that surround the alveoli.

Carbon dioxide (a waste gas from the body) passes from the blood into the alveoli, and is then breathed out.

The trachea, bronchi, and lungs are contained in the 'thoracic cavity' in the chest.

To draw air down into the thoracic cavity, the diaphragm flattens and the chest walls move out. This increases the size of the thoracic cavity, creating a negative pressure which draws air in.

Each lung is surrounded by a two layered membrane called the 'pleura'.

Between the two layers of the pleura is a thin layer of 'serous' fluid, which enables the chest walls to move freely.

The thoracic cavity is protected by the ribs, which curl around from the spine and connect to the sternum (breast bone) at the front of the body.

### What's in the air that we breathe?

| Air that we breathe in: |       | Air that we breathe out: |     |
|-------------------------|-------|--------------------------|-----|
| Oxygen                  | 20%   | Oxygen                   | 16% |
| Carbon Dioxide          | Trace | Carbon Dioxide           | 4%  |
| Nitrogen                | 79%   | Nitrogen                 | 79% |
| Other Gases             | 1%    | Other Gases              | 1%  |

### 'Normal' Respiratory Rates

|       |                          |
|-------|--------------------------|
| Adult | 12 – 20 breaths / minute |
| Child | 20 – 40 breaths / minute |
| Baby  | 30 - 60 breaths / minute |

## Choking

One of the most successful skills that can be learned by the first aider is the treatment of a casualty who is choking. Objects such as food, sweets or small objects can easily become lodged in the airway if they are accidentally 'breathed in' rather than swallowed.

### Possible signs and symptoms

- The patient is unable to speak or cough.
- Grasping or pointing to the throat.
- Distressed look on the face.
- Congestion of the face initially.
- Pale skin and cyanosis in later stages.
- Unconsciousness in later stages.

### Choking adult or child (over 1 year)

Firstly, encourage the patient to cough. If the choking is only mild, this will clear the obstruction and the patient should be able to speak to you.

If the obstruction is not cleared:

#### 1 Back blows

- Shout for help, but don't leave the patient yet.
- Bend the casualty forwards so the head is lower than the chest. For a smaller child you can place them over your knee to do this.
- Give up to 5 firm blows between the shoulder blades with the palm of your hand. Check between blows and stop if you clear the obstruction.



Back blows performed on a smaller child.



If the obstruction is still not cleared:

#### 2 Abdominal thrusts

- Stand behind the casualty (or kneel behind a small child). Place both your arms around their waist.
- Make a fist with one hand and place it just above the belly button (below the ribs) with your thumb inwards.
- Grasp this fist with your other hand, then pull sharply inwards and upwards. Do this up to 5 times. Check between thrusts and stop if you clear the obstruction.



Abdominal thrusts performed on a smaller child.



If the obstruction is still not cleared:

#### 3 Repeat steps 1 and 2

- Keep repeating steps 1 and 2.
- If the treatment seems ineffective, shout for help. Ask someone to dial 999 for an ambulance, but don't interrupt the treatment whilst the patient is still conscious.

*Abdominal thrusts can cause serious internal injuries, so send the patient to see a doctor.*

*After successful treatment, patients with a persistent cough, difficulty swallowing or with the feeling of an 'object still stuck in the throat' should also see a doctor.*

## Choking – baby (under 1 year) +

The baby may attempt to cough. If the choking is only mild, this will clear the obstruction – the baby may cry and should now be able to breathe effectively.

If the obstruction is not cleared:

### 1 Back blows

- Shout for help, but don't leave the baby yet.
- Lay the baby over your arm, face down, legs either side of your elbow with the head below the chest (see diagram).
- Give up to 5 blows between the shoulder blades with the palms of your fingers. Check between blows and stop if you clear the obstruction.



If the obstruction is still not cleared:

### 2 Chest thrusts

- Turn the baby over, chest uppermost (by laying them on your other arm) and lower the head below the level of the chest.
- Using two fingers on the chest, give up to 5 chest thrusts. These are similar to chest compressions, but sharper in nature and delivered at a slower rate. Check between thrusts and stop if you clear the obstruction.



**NEVER perform abdominal thrusts on a baby**

If the obstruction is still not cleared:

### 3 Repeat steps 1 and 2

- Keep repeating steps 1 and 2.
- If the treatment seems ineffective, shout for help. Ask someone to dial 999 for an ambulance, but don't interrupt the treatment yet.

## If the casualty becomes unconscious: +

- Support the casualty carefully to the ground (or a firm flat surface for a baby).
- **START CPR** as follows:
  - **Adult** – follow the sequence on page 7 after the heading 'if the casualty is not breathing normally'.
  - **Child** – follow the sequence on page 56 after the heading 'if the child is not breathing normally'.
  - **Baby** – follow the sequence on page 57 after the heading 'if the baby is not breathing normally'.
- Continue CPR until the child starts breathing normally on its own, help arrives to take over, or you become exhausted.



If the casualty becomes unconscious – start CPR.

## Anaphylaxis

Anaphylaxis is an extremely dangerous allergic reaction. The name 'anaphylaxis' means 'without protection' and indeed, the condition is caused by a massive over-reaction of the body's protection (*immune*) system.

Severe anaphylactic reactions are very rare. The most common reactions are to drugs (*such as penicillin*). Other common allergies are to things such as insect stings, peanuts, seafoods etc.

The main chemical that the immune cells release if they detect a 'foreign protein' is *histamine*. Histamine has several effects on the body when it is released in massive quantities:

- It makes blood vessels dilate.
- It constricts the bronchioles in the lungs.
- It makes blood capillary walls 'leaky', causing severe swelling and shock (*page 26*).
- It weakens the strength of the heart's contractions.
- It makes the skin itchy.
- It makes the skin come out in a rash.

## Possible signs and symptoms



The allergic reaction can happen in seconds, so fast recognition is essential:

- Sudden swelling of the face, tongue, lips, neck and eyes.
- Hoarse voice, 'lump in the throat', developing into loud pitched noisy breathing (*which may stop altogether*).
- Difficult, wheezy breathing, tight chest (*the patient may have the equivalent of an asthma attack as well as a swollen airway*).
- Rapid weak pulse.
- Nausea, vomiting, stomach cramps, diarrhoea.
- Itchy skin.
- Red, blotchy skin eruption.
- Anxiety – a feeling of 'impending doom'.



*This child has swelling of the tongue and lips and a red blotchy rash on his chest.*

## Treatment



- Dial 999 for an ambulance.
- Lay the casualty in a comfortable position:
  - If the casualty has Airway or Breathing problems they may prefer to sit up as this will make breathing easier.
  - **If the casualty feels faint however – do not sit them up.** Lay them down immediately. Raise the legs if they still feel faint (*page 27*).
- The casualty may carry an auto-injector of adrenaline. This can save the casualty's life if it's given promptly. The patient should be able to inject this on their own but, if necessary, assist them to use it.
- If the casualty becomes unconscious – check **Airway and Breathing** (*pages 6 to 8*) and resuscitate as necessary.
- The dose of adrenaline (*epinephrine*) can be repeated at 5 minute intervals if there is no improvement or symptoms return.



*'Epi-Pen' and 'Ana-Pen' are types of adrenaline auto-injectors.*

*Picture: many thanks to the Anaphylaxis Campaign.*

[www.anaphylaxis.org.uk](http://www.anaphylaxis.org.uk)  
© MedicalMediaKits.com



## Asthma

Asthma is a condition caused by an allergic reaction in the lungs, often to substances such as dust, traffic fumes, or pollen. Muscles surrounding the bronchioles (*see page 15*) go into spasm and constrict, making it very difficult for the patient to breathe.

Most asthma patients carry medication around with them, usually in the form of an inhaler. Ask the patient, but usually the blue inhaler is for relieving an attack, dilating the bronchioles to relieve the condition.

An asthma attack is a traumatic experience for the patient, especially a child, so reassurance and a calm approach from the First Aider is essential. If the patient is not reassured and calmed down by the First Aider, an attack can lead on to 'hyperventilation' (*see page 20*) after the inhaler has relieved the constricted airways.

### Possible signs and symptoms



- Difficulty breathing.
- Wheezy breath sounds, originating from the lungs.
- Difficulty speaking (*will need to take a breath in the middle of a sentence*).
- Pale, clammy skin.
- Grey or blue lips and skin (*cyanosis*).
- Use of muscles in the neck and upper chest to help the casualty breathe.
- Casualty will become exhausted in a severe attack.
- May become unconscious and stop breathing in a prolonged attack.



*An upright sitting position usually helps the patient to breathe more easily.*

### Treatment of asthma attack



- Help the casualty to sit upright, leaning on a table or chair if necessary.
- Help the casualty to use their reliever inhaler. This can be repeated every few minutes if the attack does not ease.
- Try to take the casualty's mind off the attack – be calm, reassuring and make light conversation.
- If the attack is prolonged, severe, appears to be getting worse, or the casualty is becoming exhausted; dial 999 for an ambulance.
- Cold winter air can make an attack worse, so don't take the casualty outside for fresh air!
- Keep the casualty upright – even if they become too weak to sit up on their own. Only lay an asthma attack patient down if they become unconscious.
- Be prepared to carry out resuscitation (*page 6 to 8*).



*Some asthma patients need to use a 'spacer device' because they can't take their inhaler all in one breath.*

## Croup

Croup is a condition usually suffered by very young children, where the larynx and trachea become infected and swollen. The attacks, which often occur during the night, can appear very alarming, but nearly always clear without causing the child any lasting harm.

### Possible signs and symptoms



- Difficult distressed breathing.
- A loud pitched, or whistling noise as the child breathes.
- A short 'barking' type cough.
- Pale, clammy skin.
- Blue tinges to the skin (*cyanosis*).
- Use of muscles in the neck and upper chest to help the child breathe.

### Treatment of croup



- Keep calm – panic will distress the child and make the attack worse.
- Sit the child up and reassure them.
- Call the doctor.
- If the attack is severe, does not ease, or the child has a temperature, dial 999 for an ambulance.

**NEVER** put your fingers down the throat of a child that appears to be suffering from croup. There is a small chance that the condition could be 'epiglottitis'. If this is the case, the epiglottis may swell even more, totally blocking the airway.

## 20 Airway and breathing problems

*“The contrasting difference between asthma and hyperventilation is the large volumes of air that can be heard entering the lungs of the hyper entilating patient, compared with the tight wheeze of the asthmatic”*

### Hyperventilation

'Hyperventilation' means 'excessive breathing'. When we breathe in, there is only a trace of carbon dioxide in the air. When we breathe out, we breathe out 4% carbon dioxide. Hyperventilating results in low levels of carbon dioxide in the blood, which causes the signs and symptoms of this condition.

A hyperventilation attack can often result from the patient being very anxious, from a panic attack or sudden fright. The condition of hyperventilation is often mistaken for 'asthma'. Asthmatics may hyperventilate after their inhalers have taken effect (*opening the airways*). The contrasting difference in the two conditions is the large volumes of air that can be heard entering the lungs of the hyperventilating patient, compared with the tight wheeze of the asthmatic.

### Possible signs and symptoms



- Unnaturally deep, fast breathing.
- Attention seeking behaviour.
- Dizziness, faintness.
- Feeling of a 'tight' chest.
- Cramps in the hands and feet.
- Flushed skin, no cyanosis.
- Pins and needles in the arms and hands.
- The patient may think they cannot breathe.
- If the attack is prolonged, the casualty may pass out and stop breathing for up to 30 seconds.

### Treatment of hyperventilation



- Be firm and calm, but reassuring with the casualty.
- Move them to a quiet place with few people around.
- Explain to the casualty that they are hyperventilating.
- 'Coach' the casualty's breathing.
- Asking the patient to take tiny sips of water will reduce the number of breaths they can take.
- Breathing through the nose will reduce the loss of carbon dioxide, but the casualty will need lots of reassurance.
- Call for medical advice if the attack is prolonged or you are in doubt.

### Drowning

Contrary to popular opinion, a casualty who drowns does not usually inhale large amounts of water into the lungs. 90% of deaths from drowning are caused by a relatively small amount of water entering the lungs, interfering with oxygen exchange in the alveoli (*wet drowning*). The other 10% are caused by muscle spasm near the epiglottis and larynx blocking the airway (*dry drowning*). The casualty will usually swallow large amounts of water, which might then be vomited as they are rescued or resuscitation takes place.

It should be remembered that other factors may contribute to the cause of drowning – for example hypothermia, alcohol, or an underlying medical condition such as epilepsy or heart attack.

### Secondary drowning

If a small amount of water enters the lungs, irritation is caused and fluid is drawn from the blood into the alveoli. This reaction could be delayed for several hours, so a casualty who has been resuscitated and 'apparently recovered' might relapse into severe difficulty breathing at a later stage. It is for this reason that any casualty who recovers from 'near drowning' should be taken to hospital immediately.

### Treatment of drowning



- Do not put yourself at risk. 'Reach or throw – don't GO'.
- If possible keep the casualty horizontal during rescue, as shock can occur.
- Check Airway and Breathing. – Perform CPR if necessary (*page 6 to 8*).
- Dial 999 for an ambulance, even if the casualty appears to recover.



**NEVER** enter the water to rescue a drowning casualty unless you have been trained to do so. Try to reach them with a rope or stick, or throw them an object that will float.  
*“Reach or throw – don't GO”*


## Collapsed lung / sucking chest wound

Each lung is surrounded by 2 layers of membrane called the 'pleura'. Between these 2 membranes is the 'pleural cavity', containing a very thin layer of 'serous fluid', which enables the two layers to move against each other as we breathe.

In a penetrating chest injury, where the outer layer of the pleura is damaged, air can be sucked from the outside of the chest into the pleural cavity, causing the lung to collapse (*pneumothorax*).

In any serious chest injury, the inner layer of the pleura could become perforated. Air may then be drawn from the lung into the pleural cavity, again causing the lung to collapse.

If air continues to be sucked into the pleural cavity, but cannot escape, pressure in the collapsed lung can build (*tension pneumothorax*). This pressure build up can squeeze the heart and the uninjured lung, making it difficult for both to function.



Collapsed lung.

## Possible signs and symptoms



- Severe difficulty breathing.
- Painful breathing.
- Fast, shallow breathing.
- Cyanosis of lips and skin.
- Pale, clammy skin.
- Uneven chest movements – the injured side of the chest may not rise.

**If there is a sucking chest wound:**

- Sound of air being drawn into the wound, with bubbling blood.
- Crackling feeling of the skin around the injury (*because of air entry*).

## Treatment



- Immediately cover a sucking chest wound with your hand (*or the casualty's hand if they are conscious*) to prevent air entry.
- Dial 999 for an ambulance – send someone to do this if possible.
- Place a sterile pad over the wound, then cover it with plastic, cling film, kitchen foil or other air tight covering.
- Tape the air tight covering on 3 sides. The dressing should prevent air from entering the wound, but still allow air to get out.
- If the casualty becomes unconscious: open the Airway, check Breathing and resuscitate if necessary. Place them in the recovery position with the injured side lowest. This will help to protect the uninjured lung.



Tape the air tight covering on 3 sides. The dressing should prevent air from entering the wound, but still allow air to get out.

## Flail chest

This is a condition where the ribs surrounding the chest have become fractured in several places, creating a 'floating' section of the chest wall.

As the casualty breathes, the rest of the chest wall moves out, but the flail segment moves inwards. As the chest wall moves back in, the flail segment moves outwards. These are called 'paradoxical' chest movements.

## Possible signs and symptoms



- Severe difficulty breathing.
- Shallow, painful breathing.
- Signs and symptoms of a fracture (*page 38*).
- 'Paradoxical' chest movements (*see above*).

## Treatment




- Dial 999 for an ambulance.
- Place the casualty in the position they find most comfortable – sat up, inclined towards the injury if possible.
- Place large amounts of padding over the flail area.
- Place the arm on the injured side in an elevated sling. Squeeze the arm gently against the padding to provide gentle, firm support to the injury.



Place padding over the flail area and place the arm on the injured side in an elevated sling.

**The circulatory system**



## The circulatory system

The circulatory system consists of a closed network of tubes (*arteries, veins and capillaries*) connected to a pump (*the heart*).


**Arteries** Carry blood away from the heart. They have strong, elastic, muscular walls which are able to expand as blood from the heart beating surges through. The largest artery, which connects directly to the heart, is called the 'aorta'.

**Veins** Carry blood towards the heart. They have thinner walls than arteries because the blood in them is under less pressure. They have one-way valves, which keep blood flowing towards the heart. The largest veins, which connect to the heart, are called 'vena cava'.

**Capillaries** Are the tiny blood vessels between the arteries and veins which allow the transfer of oxygen, carbon dioxide and nutrients in and out of the cells of the body.

**The Heart** Is a four-chambered pump. The left and right sides of the heart are separate. The left side takes blood from the lungs and pumps it around the body. The right side takes blood from the body and pumps it to the lungs.

The two sides of the heart are separated into two chambers called the 'atria' and the 'ventricles'. The **atria** are the top chambers which **collect** blood as it returns from the lungs and the body and pump it to the ventricles. The **ventricles** then pump the blood out of the heart, to the lungs and around the body.



How blood flows through the four chambers of the heart.

## The blood

60% of the blood consists of a clear yellow fluid called plasma. Suspended within the plasma are red blood cells, white blood cells, platelets and nutrients.

**Red Cells** Contain haemoglobin, which carries oxygen for use by the cells of the body. Red cells give the blood its colour.

**White Cells** Fight infection.

**Platelets** Trigger a complicated chemical reaction if a blood vessel is damaged, forming a clot.

**Nutrients** Are derived from the food by the digestive system. When combined with oxygen in the cells of the body, they provide vital energy, keeping the cell alive.

- The blood carries carbon dioxide (*the waste gas produced by the cells*) mainly in the form of 'carbonic acid'. Carbonic acid is diluted within the plasma.
- The blood also circulates heat (*generated mostly by the liver*) around the body. Heat is carried to the skin by the blood if the body needs to be cooled.



Feeling the carotid pulse.



Feeling the radial pulse.

## The pulse

Every time the heart contracts a pulsation of blood is pumped through the arteries. The walls of the arteries are elastic and expand as the blood flows rhythmically through. This expansion can be felt at the points where arteries come close to the skin.

When checking a pulse use the pads of the fingers, not the thumb (*which has its own pulse*). The First Aider should make a note of the following:

**Rate** Is it fast or slow? How many beats are there per minute?

**Rhythm** Are the beats regular? Are there any 'missed' beats?

**Strength** Does the pulse feel strong or weak?

The main pulse locations for first aid use are in the neck (*carotid pulse*), the wrist (*radial pulse*) and the upper arm (*brachial pulse*).

### Capillary refill


Circulation to the end of an arm or leg can be checked by squeezing the tip of a finger or toe. The skin will become pale when it is squeezed – if the circulation is effective, the colour should return within 2 seconds of releasing it (*this may take longer if the hands or feet are cold*).




Feeling the brachial pulse on a baby.

### 'Normal' Heart Rates at Rest


|       |                          |
|-------|--------------------------|
| Adult | 60 – 90 beats / minute   |
| Child | 90 – 110 beats / minute  |
| Baby  | 110 – 140 beats / minute |



Angina.



Heart attack.



A typical heart attack.



## Angina

Angina (*angina pectoris*) is a condition usually caused by the build up of a cholesterol plaque on the inner lining of a coronary artery. Cholesterol is a fatty chemical which is part of the outer lining of cells in the body. A cholesterol plaque is a hard, thick substance caused by deposits of cholesterol on the artery wall. Over time, the build up of the plaque causes narrowing and hardening of the artery.

During exercise or excitement, the heart requires more oxygen, but the narrowed coronary artery cannot increase the blood supply to meet this demand. As a result an area of the heart will suffer from a lack of oxygen. The patient will feel pain in the chest (*amongst other symptoms*) as a result.

Typically, an angina attack occurs with exertion, and subsides with rest. If the narrowing of the artery reaches a critical level, angina at rest (*called 'unstable angina'*) may result. A patient with angina, especially 'unstable' angina has a high risk of suffering a heart attack in the near future.

## Heart attack

Heart attack (*myocardial infarction*) is often caused when the surface of a cholesterol plaque in a coronary artery cracks and has a 'rough surface'. This can lead to the formation of a blood clot on the plaque, which completely blocks the artery resulting in the death of an area of the heart muscle.

Unlike angina, the death of the heart muscle from heart attack is permanent and will not be relieved by rest.

## Possible signs and symptoms



It should be remembered that every heart attack is different. Only a few of the signs and symptoms may be present, indeed up to a quarter of heart attacks suffered are 'silent' *without any chest pain*.

|                                 | Angina   | Heart Attack   |
|---------------------------------|--|--|
| <b>Onset</b>                    | Sudden, usually during exertion, stress or extreme weather.  | Sudden, can occur at rest.   |
| <b>Pain</b>                     | 'Vicelike' squashing pain, often described as 'dull', 'tightness' or 'pressure' on the chest. Can be mistaken for indigestion. | 'Vicelike' squashing pain, often described as 'dull', 'tightness' or 'pressure' on the chest. Can be mistaken for indigestion. |
| <b>Location of Pain</b>         | Central chest area. Can radiate into either arm ( <i>more commonly the left</i> ), the neck, jaw, back, or shoulders.          | Central chest area. Can radiate into either arm ( <i>more commonly the left</i> ), the neck, jaw, back, or shoulders.          |
| <b>Duration</b>                 | Usually lasts 3 to 8 minutes rarely longer.  | Usually lasts longer than 30 minutes.  |
| <b>Skin</b>                     | Pale, may be sweaty.   | Pale, grey colour. May sweat profusely.  |
| <b>Pulse</b>                    | Variable, depending on which area has a lack of oxygen. Often becomes irregular, missing beats.                                | Variable, depending on which area has a lack of oxygen. Often becomes irregular, missing beats.                                |
| <b>Other Signs and Symptoms</b> | Shortness of breath, weakness, anxiety.  | Shortness of breath, dizziness, nausea, vomiting. Sense of 'impending doom'.   |
| <b>Factors Giving Relief</b>    | Resting, reducing stress, taking 'G.T.N.' medication.  | 'G.T.N.' medication may give partial or no relief.   |

## Treatment of angina and heart attack

- Sit the casualty down and make them comfortable. Do not allow them to walk around. A half sitting position is often the best.
- Allow the casualty to take their own glyceryl tri-nitrate (G.T.N.) medication if they have it.
- Reassure the casualty. Remove any cause of stress or anxiety if possible.
- If you suspect heart attack – check the casualty is not allergic to aspirin, older than 16 and not already taking 'anti-coagulant' drugs (such as warfarin). If this is the case, allowing them to chew an aspirin tablet slowly may be beneficial. If you are unsure however, wait for the ambulance crew to arrive (see note, below right).

**NOTE:** Aspirin reduces the clotting ability of the blood. Chewing the tablet allows the drug to absorb quickly into the blood through the skin of the mouth, so it works faster. The ideal dose is a 300mg aspirin, but any strength will do.

- Monitor the casualty – if a heart attack victim becomes unconscious it is very likely that the heart has stopped altogether, so be prepared to perform CPR! (See pages 6 and 7).

### Dial 999 for an ambulance if:

- You suspect a heart attack.
- The casualty has not been diagnosed as having angina.
- The symptoms are different, or worse than the patients' normal angina attacks.
- Angina pain is not relieved by the patients' medication and rest after 15 minutes.
- You are in any doubt.

## Left ventricular failure

Left ventricular failure (LVF) is a condition where the left ventricle of the heart (see page 23) is not powerful enough to empty itself. The right chamber of the heart is still working properly and pumping blood into the lungs. This results in a 'back pressure' of blood in the pulmonary veins and arteries of the lungs. Fluid from this back pressure of blood seeps into the alveoli (see page 15) causing severe difficulty in breathing.

The condition can be caused by heart attack, chronic heart failure or high blood pressure. Patients with chronic heart failure often suffer attacks during the night.

## Possible signs and symptoms

- Severe difficulty in breathing.
- Crackly, often wheezy breathing (fluid on in the lungs).
- Pale sweaty skin.
- Cyanosis (blue grey tinges to skin and lips).
- Coughing frothy, blood stained sputum.
- Possibility of the signs and symptoms of heart attack.
- The patient needs to sit up to breathe.
- Anxiety, confusion, dizziness.

## Treatment

- Sit the patient up, feet dangling.
- Dial 999 for an ambulance.
- Allow the patient to take their own glyceryl tri-nitrate (G.T.N.) medication if they have it.
- Be prepared to resuscitate – the condition can quickly deteriorate.




A half sitting position is often the best.

**NOTE:** A first aider is not allowed to 'prescribe' drugs to a patient. A fully conscious adult casualty is, however, more than capable of deciding whether or not they want to take medication that may help them.



Typical G.T.N. medication that an angina patient may carry.



Normal Circulation



Hypovolaemic Shock

## Shock

To most people the word shock means an unpleasant surprise, an earthquake, or what happens if you mess about with the electrics!

The medical term shock is defined as '*inadequate tissue perfusion, caused by a fall in blood pressure or blood volume.*'

'*Inadequate tissue perfusion*' means an inadequate supply of oxygenated blood to the tissues of the body.

Now that you understand what shock is, you can understand why it can quickly result in death if not treated.

The more common causes of 'life threatening' shock are:

- **Hypovolaemic Shock**
- **Cardiogenic Shock**
- **Anaphylactic Shock**

## Hypovolaemic shock

**Hypo** means low      **vol** means volume      **aemic** means blood

This type of shock is caused by loss of body fluids, which results in a low volume of blood.

Typical causes of hypovolaemic shock are:

- External bleeding (*pages 29 and 30*).
- Internal bleeding (*page 32*).
- Burns (*pages 34 and 35*).
- Vomiting and diarrhoea (*loss of body fluids*).
- Excessive sweating.

## Possible signs and symptoms (*see also blood loss: page 29*)



The first response is release of adrenaline – this will cause:

- A rise in pulse rate.
- Pale, clammy skin (*for dark skinned casualties look at the colour of skin inside the lips*).

As the condition worsens:

- Fast, shallow breathing.
- Nausea or vomiting.
- Rapid, weak pulse.
- Dizziness, weakness.
- Cyanosis (*grey blue tinges to skin and lips*).
- Sweating.

As the brain suffers a lack of oxygen:

- Deep, sighing breathing (*air hunger*).
- Unconsciousness.
- Confusion, anxiety, even aggression.

## Treatment



- Treat the cause of the shock (*e.g. external bleeding*).
- Lay the casualty down and raise their legs in the air, returning blood to the vital organs (*take care if you suspect a fracture*).
- **Dial 999 for an ambulance.**
- Keep the casualty warm. Place a coat or blanket under the patient if they are on a cold surface, but take care not to overheat them (*as that would dilate blood vessels, causing the blood pressure to fall even more*).
- Do not allow the patient to eat, drink or smoke.
- Loosen tight clothing around the neck, chest or waist.
- Monitor breathing, pulse and levels of response.
- Be prepared to resuscitate.



Lay the casualty down and raise the legs in the air.

**NOTE:** Lay a heavily pregnant patient down leaning towards her left hand side, to prevent the baby restricting blood flow back to the heart.



## Cardiogenic shock

This is a fall in the blood pressure, caused by the heart not pumping effectively. This is the most common type of shock.


Typical causes of cardiogenic shock are:

- Heart attack (page 24).
- Cardiac failure (page 25).
- Heart valve disease.
- Tension pneumothorax (page 21).
- Cardiac arrest (page 6).

## Possible signs, symptoms and treatment



See 'Heart Conditions' (pages 24–25).



Cardiogenic Shock.

## Anaphylactic shock

Anaphylaxis is an extremely dangerous allergic reaction caused by a massive over-reaction of the body's immune system (see page 18).


An anaphylactic reaction can cause shock because the large quantity of histamine released in the body makes:

- Blood vessels dilate (causing a fall in blood pressure).
- Blood capillary walls become 'leaky' (causing a fall in blood volume).
- The strength of the heart's contractions weaker (causing a fall in blood pressure).

## Possible signs, symptoms and treatment



See 'Anaphylaxis' (page 18)



Anaphylactic Shock.

## Fainting

Fainting is caused by poor nervous control of the blood vessels and heart.

When a casualty faints, the blood vessels in the lower body dilate and the heart becomes slow. This results in the blood pressure falling and the patient has a temporary reduction in blood supply to the brain.

Typical causes of fainting are:

- Pain or fright.
- Lack of food.
- Emotional stress.
- Long periods of inactivity (such as standing or sitting).
- Heat exhaustion (page 44).

## Possible signs and symptoms




- Temporary loss of consciousness, falling to the floor.
- Slow pulse.
- Pale, clammy skin.
- Before the faint the casualty may have suffered nausea, stomach ache, blurred vision or dizziness.
- Quick recovery.

## Treatment of fainting



- Lay the casualty down and raise their legs in the air, returning blood to the vital organs.
- Check **Airway** and **Breathing** (page 6).
- Remove causes of stress, crowds of people and allow plenty of fresh air.
- Reassure the casualty as they recover. Do not allow them to sit up suddenly.
- If they feel faint again, repeat the treatment. Look for an underlying cause.
- If the casualty does not recover quickly or you are unsure: check airway and breathing again (page 6), place them in the recovery position (page 11) and dial 999 for an ambulance.



Fainting:  
The heart slows and blood vessels dilate.



Fainting – lay the casualty down and raise the legs in the air.

**Hygiene when dealing with wounds**

- *Protect yourself by covering your own cuts and abrasions with a waterproof dressing, especially on your arms and hands.*
- *Wear disposable protective gloves and an apron when you are giving first aid.*
- *Use specialised cleaning agents for cleaning up body fluid spillages. Follow the instructions on the container and use disposable towels.*
- *Dispose of soiled dressings in a yellow 'clinical waste' bag. Destroy by incineration (send the bag to hospital with the casualty if you have no clinical waste facilities).*
- *Wash your hands thoroughly before and after dealing with a patient.*
- *If you regularly deal with body fluids, ask your doctor about vaccinations against Hepatitis 'B'.*

**Wounds and bleeding**

A wound can be defined as an abnormal break in the continuity of the tissues of the body. Any wound will to some extent result in bleeding, either internally or externally. If blood loss is severe, this could result in shock (page 26), so urgent treatment would be necessary. This chapter deals with the different types of wound, the complications that may occur and their treatment.

**Types of wound and basic treatment**

- Contusion** *A bruise. Caused by ruptured capillaries bleeding under the skin. This may have been the cause of a blunt blow, or by bleeding from underlying damage, such as a fracture.*
- Cool the area with an ice pack or running water as soon as possible.
- Abrasion** *A graze. The top layers of skin are scraped off, usually as the result of a friction burn or sliding fall. Often containing particles of dirt, which could cause infection.*
- Dirt that is not embedded should be removed using clean water and sterile swabs.
  - Clean from the centre of the wound outwards, so as not to introduce more dirt into the wound.
- Laceration** *A rip or tear of the skin. More likely to have particles of dirt than a clean cut, although usually bleeds less.*
- Treat for bleeding (page 30) and prevent infection.
- Incision** *A clean cut. Usually caused by a sharp object such as a knife. Deep wounds may involve complications such as severed tendons or blood vessels. This type of wound could 'gape open' and bleed profusely.*
- Treat for bleeding (page 30) and prevent infection.
- Puncture** *A stabbing wound. Could be as a result of standing on a nail or being stabbed. The wound could be very deep and yet appear very small in diameter. Damage may be caused to underlying organs such as the heart or lungs and severe internal bleeding may occur.*
- **Dial 999 for an ambulance** if you suspect damage to underlying organs or internal bleeding.
  - Never remove an embedded object – it may be stemming bleeding and further damage may result.
- Gun Shot** *Caused by a bullet or other missile, which may be travelling at such speed as to drive into and then exit the body. A small entry wound could be accompanied by a large 'crater' exit wound. Severe bleeding and damage to organs usually results.*
- **Dial 999 for Police and Ambulance.**
  - Treat **Airway and Breathing** problems first (pages 6 to 8).
  - Pack the wound with dressings and try to prevent bleeding.
- Amputation** *Complete or partial severing of a limb.*
- See treatment of amputation (page 32).
- De-gloved** *Severing of the skin from the body, resulting in 'creasing' or a flap of skin, leaving a bare area of tissue. Caused by the force of the injuring object sliding along the length of the skin.*
- Put the skin back in place if possible.
  - Arrange urgent transport to hospital.

## Blood loss

### How much blood do we have?

The amount of blood in our body varies in relation to our size. A rough rule of thumb is that we have approximately one pint of blood per stone in body weight (*0.5 litres per 7kg*), so the average adult has between **8 and 12 pints (4.5 to 6.5 litres)** of blood, dependent on their size (*but the rule doesn't work for someone who is overweight*).

Remember that children have less blood than adults, and as such cannot afford to lose the same amount – a baby only has around **1 pint** of blood, so can only lose 1/3 of a pint before the blood pressure falls (*see below*).

### Types of bleeding

**Arterial** Blood in the arteries is under direct pressure from the heart pumping and spurts in time with the heart beat. A wound to a major artery could result in blood '**spurting**' several metres and the blood volume will rapidly reduce. Blood in the arteries is rich in oxygen and is said to be 'bright red', however this can be difficult to assess. The most important factor is *how* the wound bleeds.

**Venous** Veins are not under direct pressure from the heart, but veins carry the same volume of blood as the arteries. A wound to a major vein may '**ooze**' profusely.

**Capillary** Bleeding from capillaries occurs in all wounds. Although the flow may appear fast at first, blood loss is usually slight and is easily controlled. Bleeding from a capillary could be described as a '**trickle**' of blood.


### Effects of blood loss

The table below shows the effects, signs and symptoms of blood loss. Volumes of blood lost are given as a percentage, because we all have different amounts of blood.

As you can see, a loss of **30%** of blood volume is critical – the patient's condition rapidly deteriorates from this point onwards. Blood vessels cannot constrict any further and the heart cannot beat any faster, so blood pressure falls, resulting in unconsciousness and then death.

- Any patient with blood loss over 10% should be treated for shock (*page 26*).

See also: *Hypovolaemic shock (page 26)*  
*Hypoxia (page 14)*



|               | 10% Blood Loss   | 20% Blood Loss          | 30% Blood Loss   | 40%+ Blood Loss                            |
|---------------|--|-------------------------|--|--|
| Consciousness | Normal   | May feel dizzy stood up | Lowered levels of consciousness. Restless, anxious                         | Unresponsive                               |
| Skin          | Normal   | Pale                    | Cyanosis ( <i>blue grey tinges to the lips and skin</i> ), cold and clammy | Severe cyanosis, cold and clammy           |
| Pulse         | Normal<br><i>(this is the amount normally donated)</i> | Slightly raised         | Rapid ( <i>over 100 per min</i> ) hard to detect                           | Undetectable                               |
| Breathing     | Normal   | Slightly raised         | Rapid  | Deep sighing breaths ( <i>air hunger</i> ) |

## Treatment of external bleeding



The aims of treatment for external bleeding are firstly to stop the bleeding, preventing the casualty from going into shock (page 26), and then to prevent infection.

S.E.E.P. will help you to remember the steps of treatment:

- Sit or lay** Sit or lay the casualty down. Place them in a position that is appropriate to the location of the wound and the extent of their bleeding.
- Examine** Examine the wound. Look for foreign objects and note how the wound is bleeding. Remember what it looks like, so you can describe it to medical staff when it's covered with a bandage.
- Elevate** Elevate the wound. Ensure that the wound is above the level of the heart, using gravity to reduce the blood flow to the injury.
- Pressure** Apply direct or indirect pressure to stem bleeding:



**NEVER** try to stop bleeding by tying a band around the limb (a tourniquet) – it may cause tissue damage or make the bleeding worse.

### Direct pressure


The best way to stem bleeding is by applying direct pressure over the wound. Immediate pressure can be applied with the hands, however you should take precautions to prevent yourself from coming into contact with the patient's blood, preferably by wearing disposable gloves. The pressure should be continuous for 10 minutes. A firm bandage (*not so tight as to stop circulation to the limb altogether!*) is usually sufficient to stop bleeding from most minor wounds. If there is an embedded object in the wound, you may be able to apply pressure at either side of the object.

### Indirect pressure

If direct pressure for a wound on a limb is not possible or effective, indirect pressure can be used as a last resort. Pressure can be applied to the artery supplying the limb, squashing it against a bone and reducing the blood flow. Apply indirect pressure for a maximum of 10 minutes.

The two main indirect pressure points are:

- Brachial** Pressure is applied to the brachial artery, which runs on the inside of the upper arm. One way of doing this is to get the patient to make a fist with their opposite hand, place it under their arm pit and squeeze the injured arm down onto the fist.
- Femoral** Pressure is applied to the femoral artery, which is located where the thigh bone (*femur*) crosses the 'bikini line'. Take care to explain your actions. One way of doing this is to use the heel of your foot to apply the pressure.



Indirect Pressure Points

### Dressings

A dressing should be sterile and just large enough to cover the wound. It should be absorbent and preferably made of material that won't stick to the clotting blood (*a 'non-adherent' dressing*).

A firmly applied dressing is sufficient to stem bleeding from the majority of minor wounds, but the dressing should not restrict blood flow to the rest of the limb (*check the circulation with a 'capillary refill' test, page 23*)

Extra pressure 'by hand' and elevation may be necessary for severe bleeding. If the dressing becomes saturated with blood, keep it in place and put another larger dressing on top. If this doesn't work take the dressings off and start again.



## Embedded objects



### Objects embedded in a wound:

An object embedded in a wound (other than a small splinter) should not be removed as it may be stemming bleeding, or further damage may result.

Use sterile dressings and bandages to 'build up' around the object. This will apply pressure around the wound and support the object. Send the casualty to hospital to have the object removed.

### Splinters:

If a splinter is embedded deeply, difficult to remove or on a joint, leave it in place and follow the advice for embedded objects above. Other splinters can be removed as follows:

- Carefully clean the area with warm soapy water.
- Using a pair of clean tweezers, grip the splinter as close to the skin as possible. Gently pull the splinter out at the same angle that it entered.
- Gently squeeze around the wound to encourage a little bleeding. Wash the wound again, then dry and cover with a dressing.
- Seek medical advice to ensure the casualty's tetanus immunisation is up-to-date.

### Objects embedded in the nose, ear or other orifice:

Do not attempt to remove anything that someone has got stuck in their ear, nose or other orifice. Take them to hospital where the professionals can remove it safely.



## Nose bleeds



Weakened or dried out blood vessels in the nose can rupture as a result of a bang to the nose, picking or blowing it. More serious causes could be high blood pressure or a fractured skull.

- Sit the patient down, head tipped forward.
- Nip the soft part of the nose. Maintain constant pressure for 10 minutes.
- Tell the patient to breathe through the mouth.
- Give the patient a cloth to mop up any blood whilst the nose is nipped.
- Advise the patient not to breathe through or blow their nose for a few hours after bleeding has stopped.
- If bleeding persists for more than 30 minutes, or if the patient takes 'anti-coagulant' drugs (such as warfarin), take or send them to hospital in an upright position.
- Advise a patient suffering from frequent nosebleeds to visit their doctor.



*Nip the soft part of the nose. Maintain constant pressure for 10 minutes.*

## Eye injury



Small particles of dust or dirt can be washed out of an eye with cold tap water. Ensure the water runs away from the good eye.

### For a more serious eye injury:

- Keep the casualty still and gently hold a soft sterile dressing over the injured eye. This can be carefully bandaged in place if necessary.
- Tell the casualty to close their good eye, because any movement of this will cause the injured eye to move also. If necessary bandage the good eye to stop the casualty using it. Lots of reassurance will be needed!
- Take the casualty to hospital. Dial 999 for an ambulance if necessary.

### For chemicals in the eye:

- Wear protective gloves. Wash with copious amounts of clean water, ensuring the water runs away from the good eye. Gently but firmly try to open the casualty's eyelid to irrigate the eye fully. Dial 999 for an ambulance.





## Amputation



Amputation is the complete or partial severing of a limb, and is extremely traumatic for the casualty. Your priorities are to stop any bleeding, to carefully preserve the amputated body part and to reassure the casualty.

- Treat the casualty for bleeding (*page 30*) and for shock (*page 26*).
- Dial 999 for an ambulance.
- Dress the casualty's wound with a 'low-adherent', non-fluffy dressing.
- Wrap the amputated part in a plastic bag, and then put the package on a bag of ice to preserve it.  
**Do not allow the amputated part to come into direct contact with the ice or get wet.**

## Crush injury

Crush injuries most commonly occur as a result of building site or traffic accidents. If the blood flow to a limb (*e.g. an arm or a leg*) is impaired by the weight of a crushing object, there is a danger of toxins building up in the muscle tissues below the site of the crushing.

If the blood flow to the limb is impaired for 15 minutes or more, the toxins will build up so much that if they are released into the rest of the body (*which will happen when the crushing object is removed*) they may cause kidney failure. This is called 'crush syndrome' and may result in death.

Expert medical care is needed when releasing the patient if the blood flow has been impaired for 15 minutes or more.

### Treatment for crushing less than 15 minutes



- Release the casualty as quickly as possible if you can.
- Dial 999 for an ambulance.
- Control any bleeding and cover open wounds.
- Treat for shock if necessary (*page 26*), taking care not to move injuries.
- Monitor Airway and Breathing until help arrives.

### Treatment for crushing more than 15 minutes



- DO NOT release the casualty.
- Dial 999 for an ambulance. Give clear information about the incident.
- Monitor Airway and Breathing until help arrives.

## Internal bleeding

Internal bleeding is a very serious condition, yet can be very difficult to recognise in its early stages. Internal bleeding can be as a result of injury, such as lung or abdominal injuries, yet can also happen 'spontaneously' to an apparently well patient, such as bleeding from a stomach ulcer or a weak artery.

Although blood may not actually be lost 'externally' from the body, it is lost out of the arteries and veins, so shock can quickly develop.

Other serious life threatening complications can occur from internal bleeding, such as a brain haemorrhage or bleeding into the lungs.

### Possible signs and symptoms



You should suspect internal bleeding if signs of shock (*see pages 26 and 29*) are present, but there is no obvious cause, such as external bleeding.

There may be:

- Signs of SHOCK (*page 26*).
- Bruising and/or swelling.
- Pain, or a history of recent pain at the site of bleeding.
- Other symptoms related to the site of bleeding (*e.g. difficulty breathing if the bleeding is in the lung*).

### Treatment of internal bleeding



- Dial 999 for an ambulance.
- Treat the casualty for shock as necessary (*page 26*).

## Poisons

A poison can be described as any substance (*solid, liquid or gas*) that causes damage when it enters the body in a sufficient quantity.

Poisons can enter the body in 4 ways, they can be:

- Ingested** Swallowed, either accidentally or on purpose.
- Inhaled** Breathed in, accessing the blood stream very quickly as it passes through the alveoli.
- Absorbed** Through the skin (*see chemical burns, page 35*).
- Injected** Through the skin, directly into tissues or a blood vessel.

A poison can either be:

- Corrosive** Such as: acids, bleach, ammonia, petrol, turpentine, dishwasher powder, etc.

OR

- Non-Corrosive** Such as: tablets, drugs, alcohol, plants, perfume etc.



## Possible signs and symptoms



The signs and symptoms of poisoning are wide, varied and dependent on the substance. Look for clues such as:

- Containers or bottles.
- Syringe or drug taking equipment.
- Tablets or drugs.
- Smell on the breath.

### Other signs that can accompany poisoning may be:

- Vomiting or retching.
- Confusion or hallucination.
- Abdominal pains.
- Headache.
- Burns (*or burning sensation*) around the entry area.
- Unconsciousness, sometimes fitting.
- Breathing problems.
- Cyanosis.

## Treatment



### For a corrosive substance:

- Don't endanger yourself – make sure it's safe to help.
- Dilute the substance or wash it away if possible:
  - Substances on the skin – see chemical burns (*page 35*).
  - Ingested Substances – get the casualty to rinse out their mouth, then give frequent sips of milk or water.
- **Dial 999 for an ambulance.** Give information about the poison if possible. Take advice from the ambulance operator.
- If the casualty becomes unconscious – open the **Airway** and check for **Breathing**. Resuscitate as necessary using a protective face-shield (*pages 6 to 8*). If the casualty is breathing effectively, place them in the recovery position, then **dial 999 for an ambulance**.

### For a non-corrosive substance:

- **Dial 999 for an ambulance.** Give information about the poison if possible. Take advice from the ambulance operator.
- If the casualty becomes unconscious – open the **Airway** and check for **Breathing**. Resuscitate as necessary using a protective face-shield (*pages 6 to 8*). If the casualty is breathing effectively, place them in the recovery position, then **dial 999 for an ambulance**.



*Get the casualty to rinse out their mouth, then give frequent sips of milk or water.*

**NEVER** make the patient vomit. This may put the airway in danger.

*It helps the Paramedics if you:*

- *Pass on containers, or other information about the substance.*
- *Find out how much has been taken.*
- *Find out when it was taken.*
- *Keep samples of any vomit for hospital analysis.*

## Burns and scalds

### Estimating the severity of a burn



There are 5 factors that combine to affect the severity of a burn:

- Size** The larger the area of the burn, the more severe. The size of the burn is given as a percentage of the body's surface area. An easy way to work this out is to compare the size of the burn with the patient's hand. An area equal to the size of the palm of the patient's opened hand (*including fingers*) is equal to 1% of their body area.
- Cause** The cause of the burn, as previously described in this chapter, will influence the overall severity – for example, electrical burns may leave a patient with deep internal burns. Some chemicals (*such as hydrofluoric acid*) could cause poisoning in addition to burns.
- Age** The age of the patient will affect the recovery rate and severity. Babies and young children will burn at lower temperatures than adults. Elderly patient's burns take longer to heal and they may be more susceptible to infection.
- Location** The location of the burn can affect the severity – in particular burns to the airway of a patient by inhaling hot gasses can be an instant killer. Burns to the eye may result in blindness.
- Depth** The deeper the burn, the more severe. *See depth of burns below.*



An area equal to the size of the palm of the patient's opened hand (*including fingers*) is equal to 1% of their body area.

### Depth of burns



The skin consists of 3 layers – the 'epidermis' on the outside, the 'dermis' beneath, which lies on a layer of 'subcutaneous' fat.

The depth of burns can be defined as:

- Superficial** This involves only the outer epidermis layer, and most commonly occurs from scalds. The burn looks red, sore and swollen.
- Intermediate** This affects both the epidermis and the dermis layers of skin. The burn looks raw and blisters will form.
- Full Thickness** The layers of skin are burned away to the subcutaneous fat layer or beyond. The burn may look pale, charred or waxy. The nerve endings will be burned away, so pain in this area may be absent, misleading both you and the patient.

## Causes of burns and treatment



The different causes of burn can be separated in to 5 areas. The treatment for the burn can differ slightly depending on the cause:

### Electric burns

Caused by heat that is generated by an electrical current flowing through the tissues of the body. You may be able to see a burn where the current entered the body, and at the point of exit. There may be deep internal burns which are not visible along the path of the current flow. The extent of the internal burns can be estimated by the severity of the entry and exit wounds.

An electric shock may cause cardiac arrest. In this case, Airway and Breathing become the priority.

- Ensure your own safety – make sure contact with the electricity is broken.
- Ensure Airway and Breathing are maintained (*pages 6 to 8*).
- Irrigate the area of the burns, including the path between entry and exit, for at least 10 minutes.
- Dial 999 for an ambulance.
- Continue treatment as you would for a 'dry heat' burn.

**NEVER** burst blisters (*the layer of skin is protecting against infection*)

**NEVER** touch the burn.

**NEVER** apply lotions, ointments or fats – they might introduce infection, and would need to be removed in hospital.

**NEVER** apply adhesive tape or dressings – the burn may be larger than it first appears.

**NEVER** remove clothing that has stuck to the burn.



## Dry heat burns

Any direct contact with a dry heat source or friction.

- Do not put yourself in danger.
- Ensure that Airway and Breathing are maintained (*page 6*).
- **Cool** the burn immediately with cold (*preferably running*) water, for at least 10 minutes. If water is not available, any cold harmless liquid (*e.g. milk*) is better than no cooling at all. Do this first then move quickly to a water supply if you can. Take care not to cool large areas of burns so much that you induce hypothermia.
- **Remove** watches, rings etc. during cooling, as burned areas will swell. Clothing that has not stuck to the burn may be removed very carefully.
- **Dress** the burn with a sterile dressing that won't stick. Cling film is one of the best dressings for a burn – discard the first two turns from the roll and apply it lengthways (*don't wrap it tightly around a limb*). Secure with a bandage.
- Alternative dressings could be a new, unused plastic bag, low adherent dressings or specialised burns dressings (*do not rely on burns dressings to cool a burn – use cold water*).
- See note (*below right*) on when to seek medical advice.
- **Dial 999 for an ambulance** if the burn appears severe, or the casualty has breathed in smoke or fumes.



Cool the burn for 10 minutes.



Remove jewellery and LOOSE clothing.



Dress the burn. Cling film is one of the best dressings for a burn.

## Wet heat (scalds)

Scalds are most commonly from hot water, but may be from hot fats or other liquids that can reach higher temperatures than water.

- Treat as a dry heat burn.

## Chemical burns

Caused by chemicals which either corrode the skin or create heat (*or both*).

It is important to learn the correct first aid treatments for any chemicals used in your workplace – different chemicals can have different first aid treatments.

- Make the area safe – contain the chemical if possible and protect yourself from coming into contact with it.
- Dry powder chemicals can be carefully brushed off the skin before irrigating. Take care to protect yourself.
- Irrigate the burn with lots of running water to wash the chemical away. This should be done for longer than a thermal burn – at least 20 minutes. Take care not to wash the chemical onto unaffected areas of the body. Ensure pools of contaminated water do not collect underneath the casualty.
- **Dial 999 for an ambulance.** Make a note of the chemical, and give this information to the ambulance operator if possible.
- Remove contaminated clothing carefully whilst irrigating the burn.
- If an eye is contaminated, irrigate as above, and ensure that the water runs away from the un-affected eye (*see page 31*).
- Some chemicals in the workplace cannot be safely diluted with water – health and safety regulations require an 'antidote' to be available in an emergency. You should be trained in the use of the antidote.

## Radiation burns (sun burn)

Most commonly seen as sunburn.

- Remove the casualty from exposure to the sun; indoors if possible.
- Give the casualty frequent sips of water to ensure that heat exhaustion does not take effect (*page 44*).
- Cool the burn with cold water. If the area affected is extensive, cool the burn under a gentle cold shower or in a bath of cold water for 10 minutes.
- If there is extensive blistering, or you are not sure, seek medical advice.
- If the sunburn is mild, after-sun cream or calamine lotion may soothe the area.

### Seek medical advice if:


- The burn is larger than 1-inch square.
- The patient is a child.
- The burn goes all the way around a limb.
- Any part of the burn appears to be full thickness.
- The burn involves hands, feet, genitals or the face.
- You are not sure.

## 36 Injuries to bone, muscles and joints

The skeleton consists of 206 bones, the functions of which are to:

- Provide support for the soft tissues of the body. This gives the body its shape.
- Provide protection for important organs such as the brain, lungs and spinal cord.
- Allow movement, by incorporating different types of joints and attachment for muscles.
- Produce red blood cells, some white blood cells and platelets in the marrow of bones such as the femur.
- Provide a store of minerals and energy such as calcium and fats.

### The skeletal system



## Causes of injury


Injury can be caused to the bones, muscles and joints by different types of force:

- Direct Force**     Damage results at the location where the force was applied, e.g. as the result of a blow or kick.
- Indirect Force**     Damage occurs away from the point where the force was applied, e.g. a fractured collar bone, as a result of landing on an outstretched arm.
- Twisting Force**     Damage results from torsion forces on the bones and muscles, Force e.g. 'twisting an ankle'.
- Violent Movement**     Injury results from a sudden violent movement, such as injuring the knee joint by kicking violently.
- Pathological**     Injury results because the bones have become brittle or weak, due to disease or old age.


## Types of fracture

A fracture can be defined as a 'break in the continuity of the bone'. The basic categories of fracture are:


- Closed**     This is a clean break or crack in the bone, with no complications.
- Open**     The skin has become broken by the bone which may (*or may not*) still be protruding from the wound. This type of injury has a high risk of infection.
- Complicated**     With this type of injury, there are complications which have arisen as a result of the fracture, such as trapped blood vessels or nerves.
- Green Stick**     This type of fracture occurs more commonly in children, who have young, more flexible bone. The bone is split, but not totally severed. Green Stick fractures are often mistaken for sprains and strains, because only a few of the signs and symptoms of a fracture are present.




*Closed Fracture*



*Open Fracture*



*Complicated Fracture*



*Green Stick Fracture*

## Dislocations

A dislocation is where a bone becomes partially or fully dislodged at a joint, usually as a result of wrenching movement or sudden muscular contraction. The most common dislocations are the knee cap, shoulder, jaw, thumb or a finger.

There may also be a fracture at or near the site of the dislocation, and damage to ligaments, tendons and cartilage. It can be difficult to distinguish between a fracture and a dislocation.

Never attempt to manipulate a dislocated joint back into place. This is a job for the experts – the procedure can be extremely painful for the patient, and you may cause further damage.

## Sprains and strains

A *sprain* is defined as an injury to a ligament at a joint. A *strain* is defined as an injury to muscle. Usually caused by sudden wrenching movements, the joint overstretches, tearing the surrounding muscle or ligament.

Minor fractures are commonly mistaken for sprains and strains. If you are not sure, you should treat the injury as if it was a fracture. The only way to rule out a fracture is by x-ray.

For treatment of sprains and strains – see page 39

## Support sling

1



2



3



## Elevated sling

1



2



3



## Possible signs and symptoms of a fracture



|                             |  |
|-----------------------------|--|
| <b>Pain</b>                 | At the site of the fracture. Strong pain killers, nerve damage or dementia may mask the pain, so beware.     |
| <b>Loss of Power</b>        | e.g. not being able to lift anything with a fractured arm.   |
| <b>Unnatural movement</b>   | This type of fracture is classed as 'unstable' and care should be taken to prevent the fracture from moving. |
| <b>Swelling or bruising</b> | Around the site of the fracture.   |
| <b>Deformity</b>            | If a leg is bent in the wrong place, it's broken!  |
| <b>Irregularity</b>         | Lumps or depressions along the surface of the bone, where the broken ends of the bone overlap.               |
| <b>Crepitus</b>             | The feeling and sound of bone grating on bone, when the broken ends rub on each other.                       |
| <b>Tenderness</b>           | At the site of the injury.   |

## Treatment of a basic fracture



See also: **Head Injuries** (pages 12 to 13)  
**Flail Chest** (page 21)  
**Spinal Injuries** (pages 39 to 40)

- Reassure the casualty, tell them to keep still.
- Keep injury still with your hands until it is properly immobilised. The casualty might be able to do this on their own.
- Don't move the casualty until the injury is immobilised, unless they are in danger.
- Don't try to bandage an injury if you have called an ambulance, just keep it still (*cover open wounds with a sterile dressing*).
- Don't let the casualty eat or drink – they may need an operation.

## For an upper limb injury:

- Carefully place the arm in a sling against the trunk of the body. Arm fractures are normally placed in a support sling. Collar bone fractures are normally supported by an elevated sling (*keep the elbow down at the patient's side when using an elevated sling for a collar bone fracture*).
- If the casualty is in severe pain, circulation or nerves to the arm are affected, the casualty has breathing difficulties, or you are unsure, **dial 999 for an ambulance**.
- Arrange transport to hospital.

## For a lower limb injury:

- Keep the casualty warm and still. **Dial 999 for an ambulance**.
- If the ambulance arrival will be delayed (*e.g. remote countryside*) immobilise the injury by bandaging the sound leg to the injured one.
- Check circulation beyond the injury and any bandages. Loosen bandages if necessary.

## Treatment of sprains and strains



The best treatment for a sprain or strain is to follow the RICE mnemonic:

- R**est                 Rest the injury. e.g. don't allow a sports player to carry on playing (*it's better to take time out now than miss the next ten matches!*).
- I**ce                    Apply an ice pack to the injury as soon as possible. This will help reduce swelling, which will speed recovery. Place a tea towel or triangular bandage between the skin and the ice pack. Do this for 10 minutes, every 2 hours, for 24 hours for maximum effect.
- C**ompression       Apply a firm (*not constrictive*) bandage to the injured area. This helps to reduce swelling. The bandage can be applied over a crushed ice pack for the first 10 minutes.
- E**levation            Elevate the injury. This also reduces swelling.

Remember: minor fractures can easily be mistaken for sprains and strains. The only way to rule out a fracture is by x-ray, so take or send the casualty to hospital.


**CAUTION:** To prevent frostbite always wrap the ice pack in a cloth and apply it for a maximum of 10 minutes. Allow the skin to return to normal temperature before repeat applications.



## Spinal injuries


Spinal injury occurs with approximately 2% of trauma (*injury*) patients. Although this figure appears relatively low, suspecting and correctly treating the injury is essential, because poor treatment of a patient with a spinal injury could result in them becoming crippled for life or even death.

The spinal cord is an extension of the brain stem, and travels down the back of the spinal vertebrae. Vital nerves, controlling breathing and movement of limbs travel down the spinal cord (*see diagram*). The weakest part of the spinal column is the neck, and indeed a neck injury can be the most severe type of spinal injury, because the nerves controlling breathing may become severed.




## Suspect spinal injury if the casualty has:

- Sustained a blow to the head, neck or back (*especially resulting in unconsciousness*).
- Fallen from a height (*e.g. fall from a horse*).
- Dived into shallow water.
- Been in an accident involving speed (*e.g. car accident or knocked down*).
- Been involved in a 'cave in' accident (*e.g. crushing, or collapsed rugby scrum*).
- Multiple injuries.
- Pain or tenderness in the neck or back after an accident (*pain killers or other severe injuries may mask the pain – beware*).
- OR: if you are in any doubt.




The spinal cord travels through the centre of the spinal column. Nerves emanate from each vertebrae in pairs.

Possible signs and symptoms of spinal injury 

Remember – If some of these signs and symptoms are present, nerves may already be damaged. You should treat a patient who you *suspect* has a spinal injury to *prevent* these signs and symptoms from developing.

- Pain or tenderness in the neck or back.
- Signs of a fracture in the neck or back (*page 38*).
- Loss of control of limbs at or below the site of injury.
- Loss of feeling in the limbs.
- Sensations in the limbs, such as pins and needles or burning.
- Breathing difficulties.
- Incontinence.

Treatment of spinal injury 

## If the patient is conscious:

- Reassure the patient. Tell them not to move.
- Keep the patient in the position you find them. Do not allow them to move, unless they are in severe danger.
- Hold their head still with your hands. Keep the head and neck in line with the upper body (*see diagram*).
- Dial 999 for an ambulance. Keep the patient still and warm until it arrives.

## If the patient is unconscious and breathing normally:

- Do not move the patient unless they are in severe danger.
- If the patient is breathing normally this means the airway must be clear, so there is no need to tip the head back. The 'jaw thrust' technique can be used to keep the airway open without moving the head (*this is explained on page 41*). Constantly monitor breathing.
- Dial 999 for an ambulance.
- Hold the head still with your hands. Keep the head and neck in line with the upper body (*see diagram*).
- If you have to leave the casualty, if they begin to vomit, or if you are concerned about their airway in any way, place the casualty in the recovery position. Keep the head, neck and upper body in line as you turn the patient. Doing this effectively takes more than one rescuer, so get local help if you can (*see page 41 for methods of turning a spinal injury patient*).
- Keep the casualty warm and still. Constantly monitor Airway and Breathing until help arrives (*page 6*).

## If the patient is not breathing normally:

- If the patient is not breathing normally, the airway will need to be opened. Head tilt may be used, but the tilt should be the minimum that is required to allow unobstructed rescue breaths.
- Only if you are trained and confident, you can try the 'jaw thrust' technique to open the airway, but if you find the patient is still not breathing normally, you should then open the airway using the head tilt method before carrying out resuscitation (*page 6*).
- Re-check breathing once the airway has been opened.
- If the casualty is still not breathing normally, dial 999 for an ambulance, then carry out resuscitation (*pages 6 to 8*).
- Obtain the help of others to support the head as you resuscitate.

Remember – successful resuscitation that results in paralysis from a neck injury is a tragedy, but failing to maintain an adequate airway will result in death.



Holding a patient's head still in a car.



Keep the head, neck and upper body in line.

## Managing the airway with spinal injuries +

If a patient is unconscious and laid on their back, the airway can be in danger from vomit or the tongue falling back.

A patient who has not been injured can simply be turned into the recovery position to protect the airway, but if spinal injuries are suspected, great care must be taken not to move the spine.

If a patient is already on their side (*not on their back*) you may not have to move them at all. Is the airway in danger from vomit or the tongue falling back? If not, the patient can be kept still in the position you find them.

If you can continually monitor that the patient is breathing normally, you may be able to keep them still until the ambulance arrives, even if they are on their back.

If the tongue begins to fall back or the patient vomits however, immediate action will be needed to protect the airway:

### Jaw thrust

If the patient is breathing but the tongue is starting to obstruct the airway (*usually makes snoring type noises*) the jaw thrust technique can be used to keep the airway open:

- Kneel above the head of the patient, knees apart to give you balance.
- With your elbows resting on your legs (*or the floor*) for support, hold the patient's head with your hands to keep their head and neck in line with the body (*see fig.1*).
- Place your middle and index fingers under the jaw line of the patient (*under their ears*).
- Keeping the head still, lift the jaw upwards with your fingers (*see fig.2*). This gently lifts the tongue from the back of the throat.

**DO NOT** attempt the jaw thrust technique during CPR – tilt the head to open the airway instead (*page 6*).

### Log roll

If you have to leave the casualty, if they begin to vomit, or if you are concerned about their airway in any way, the patient will have to be turned onto their side. The head, neck and upper body must be kept in line as you turn the patient.

The best method of turning a spinal injury patient is the log roll technique, but you will need at least three helpers to roll the patient.

- Support the head of the patient, keeping the head, neck and upper body in line (*see fig.1*).
- Your helpers should kneel along one side of the patient. Get them to gently straighten the patient's legs and arms.
- Making sure that everyone works together, the helpers should roll the patient towards them on your count. You gently move the head to follow the body as the patient is rolled. (*see fig.3*).
- Keep the head, neck, body and legs in line at all times. If you can, keep the patient in this position until the ambulance arrives.

### Recovery position

If the patient has to be turned onto their side and you don't have three helpers, you will need to use the recovery position method when turning the patient. Keep the head, neck and body in line as best as you can as you roll the patient over. Have some padding (*e.g. a folded coat*) to support the patient's head when they are on their side.

If you have one or two helpers, you can support the head as your helper(s) turn the patient.

- Start by supporting the head of the patient, keeping the head, neck and upper body in line (*see fig.1*).
- Get your helper(s) to gently move the patient's arms and legs into position, ready to turn the patient into the recovery position (*see fig.4*).
- Making sure that everyone works together, the helper(s) should roll the patient into the recovery position. The helper(s) should pull equally on the patient's far leg and shoulder as they turn the patient, keeping the spine in line. You gently move the head to keep it in line with the body as the patient is moved (*see fig.5*).



Fig. 1: Keeping the patient's head and neck in line with the body.



Fig. 2: The jaw thrust technique. Use your middle and index fingers to lift the jaw whilst you keep the head still.



Fig. 3: Log roll.



Fig. 4: Get your helper(s) to position the patient's arm and legs ready for the recovery position.



Fig. 5: Get your helper(s) to turn the patient whilst you keep the head in line with the body.

## 42 Effects of heat and cold

This chapter covers the effects of over exposure to heat or cold on the body.

Severe Hypothermia or Heat Stroke are potentially fatal conditions, and need skilful treatment from the First Aider.

The people who are most at risk from the effects of heat and cold are the elderly or infirm, babies and children, or people who take part in outdoor activities such as hiking or sailing.

### Body temperature

The body works best when its temperature is close to 37°C (98.6°F). This temperature is maintained by an area in the centre of the brain called the 'hypothalamus'.

If the body becomes too hot we produce sweat, which evaporates and cools the skin. Blood vessels near to the skin dilate (*flushed skin*) and the cooled blood is circulated around the body.


If the body becomes too cold we shiver, which creates heat by muscle movement. Blood vessels near to the skin constrict (*pale skin*), keeping the blood close to the warmer core of the body. Hairs on the skin become erect, trapping warm air (*goose pimples*).

Injuries resulting from exposure to extremes of temperature can be 'localised' (such as *sunburn or frostbite*), or 'generalised' (such as *hypothermia or heat stroke*).

### Signs and symptoms of body temperature change



The symptoms of over-exposure to heat or cold are demonstrated by the diagram below. As the temperature of the body becomes too hot or too cold, the area of the brain that regulates temperature (*the hypothalamus*) stops working, and the condition rapidly becomes worse as the body no longer fights the condition:



See Also:  
Taking a temperature (page 45)



## Hypothermia

The onset of hypothermia occurs when the body's core temperature falls below 35°C. A patient suffering hypothermia in its mildest form who is treated effectively will usually make a full recovery. If the body's core temperature falls below 26°C the condition will most likely be fatal, however resuscitation has been successful on people with temperatures as low as 10°C, so it is always worth attempting.

The underlying cause of hypothermia is over exposure to cold temperatures, however different conditions and types of patient will increase the risk:

- The hypothalamus (*temperature control centre*) of a baby or young child is under developed, and hypothermia can result from as little as being in a cold room.
- Elderly or infirm patients do not generate as much body heat, so prolonged periods in a cold environment can lower the core temperature.
- Wet clothing, or immersion in cold water results in the body cooling much faster than it would in dry air. Water conducts heat away from the body.
- A person who is not clothed properly in windy conditions will have cold air continually in contact with the skin, resulting in faster cooling of the body.



## Possible signs and symptoms



- Pale skin, cold to touch.
- Shivering at first, then muscle stiffness as the body cools further.
- Slowing of the body's functions – including thought, speech, pulse and breathing (*the pulse can fall lower than 40 beats per minute*).
- Lethargy, confusion, disorientation (*can be mistaken for drunkenness*).
- Lowered levels of response, eventually unconsciousness, then death.

## Treatment



### If the casualty is unconscious:

- Open the Airway and check Breathing. Resuscitate if necessary (*pages 6 to 8*).
- Dial 999 for an ambulance.
- Gently place the patient in the recovery position (*page 11*). Do not move the patient unnecessarily, because the slightest jolt can stop the heart.
- Place blankets or other insulating materials under and around the patient. Cover the head.
- Constantly monitor breathing. The pulse may be hard to find – it is safe to assume the heart is beating if the casualty is breathing normally.

### For a conscious casualty:

- If you can shelter the casualty, remove any wet clothing. Quickly replace with dry, warm garments. Cover the head.
- If the casualty is fit, young and able to climb into a bath without help, bathe them in warm water (40°C / 104°F). Don't allow an elderly patient to bathe.
- If a bath is not possible, wrap them in warm blankets. Heat the room to a warm temperature (25°C / 77°F) if indoors.
- A casualty outdoors should be insulated from the environment and ground. Use a survival bag and shelter if available. Share your body heat with them.
- Give the casualty warm drinks and food.
- Seek medical advice if the patient is elderly, a child, or if you are in any doubt about their condition.
- If the condition seems severe. Dial 999 for an ambulance.

**NEVER** give a patient alcohol (*it dilates blood vessels, which will make the patient colder*).

**NEVER** place direct sources of heat on or near the patient (*they draw blood to the skin, causing a fall in blood pressure and place stress on the heart*).

**NEVER** warm babies or the elderly too quickly (*e.g. by placing them in a warm bath*).

**BEWARE:** A hypothermic heart is in grave risk of 'ventricular fibrillation', which causes cardiac arrest. Handle hypothermic patients with care – the slightest jolt can induce the condition.

## 44 Effects of heat and cold

### Trench Foot

This is caused by prolonged exposure to wet, cold conditions. The cells do not freeze, so full recovery is usual. The symptoms and treatment are similar to frostbite.

### Chilblains

The most common cold injury, caused by exposure to dry cold. Again the cells do not freeze. There may be itching, reddish-blue skin and swelling. With time, blisters may form. Treat as frostbite.

**NEVER** rub the affected area.

**NEVER** use direct or dry heat to warm the injury.

**NEVER** re-warm the injury if there is a danger of it refreezing.



Give the casualty drinks of water to re-hydrate them.

## Frostbite

Frostbite is a condition caused when an extremity (such as a finger or an ear) is subject to cold conditions. The cells of the limb become frozen. Ice crystals form in the cells, which causes them to rupture and die. Frostbite may also be accompanied by hypothermia, which should also be treated. Serious frostbite can result in the complete loss of a limb, particularly fingers or toes.

### Possible signs and symptoms

- Pins and needles, followed by numbness.
- Hardening and stiffening of the skin.
- Skin colour change – first white, then blue tinges, then eventually black.
- On recovery, the injury will become hot, red, blistered and very painful.

### Treatment

- Gently remove rings, watches etc.
- Stop the freezing becoming worse if the casualty is still outdoors – place the limb under their arm or hold it with your hands.
- Don't rub the injury – this will cause damage.
- Don't re-warm the injury if there is a risk of it refreezing. Move the patient indoors before you treat them.
- Place the injury in warm water (test the temperature with your elbow as you would for a baby's bath – not with a frozen hand!).
- An adult casualty can take two paracetamol tablets for intense pain.
- Take the casualty to hospital as soon as possible.

## Heat exhaustion

Heat exhaustion is the body's response to loss of water and salt through excessive sweating. The most common cause of this condition is working or exercising in hot conditions (such as hiking on a very hot day).

Heat exhaustion occurs when the core body temperature raises above 38°C. If the problem is not treated, it can quickly lead to heat stroke (opposite).

### Possible signs and symptoms

- Confusion, dizziness.
- Pale, sweaty skin.
- Nausea, loss of appetite, vomiting.
- Fast, weak pulse and breathing.
- Cramps in the arms, legs, abdomen.
- The casualty may say that they 'feel cold', but they will be hot to touch.

### Treatment

- Take the casualty to a cool place.
- Remove excessive clothing and lay them down.
- Give the casualty plenty of water to re-hydrate them. Oral rehydration solutions (such as 'Dioralyte') or isotonic drinks are best as they also replace lost salt.
- Obtain medical advice, even if the casualty recovers quickly.
- If the casualty's levels of response (page 9) deteriorate – place them in the recovery position and dial 999 for an ambulance. Monitor Airway and Breathing (page 6).
- Treat for heat stroke (opposite) as necessary.

## Heat stroke

Heat stroke is a very serious condition. It results from failure of the hypothalamus (*temperature control centre*) in the brain. The sweating mechanism fails, the body is unable to cool down and the core temperature can reach dangerously high levels (*over 40°C*) within 10 to 15 minutes.

The condition can be caused by a high fever or prolonged exposure to heat and often follows heat exhaustion (*previous page*).

## Possible signs and symptoms



- Severe confusion and restlessness.
- Flushed, hot, dry skin (*no sweating*).
- Strong, fast pulse.
- Throbbing headache.
- Dizziness.
- Nausea, vomiting.
- Reduction in levels of response (*page 9*) leading to unconsciousness.
- Possibility of seizures if unconscious.

## Treatment



- Move the casualty to a cool, shaded area.
- **Dial 999 for an ambulance.**
- Cool the casualty rapidly, using whatever methods you can:
  - Remove outer clothing, and wrap the casualty in a cold, wet sheet. Keep it wet and cold until the casualty's temperature falls to normal levels, then replace with a dry sheet.
- Other methods of cooling can be:
  - Continually sponging with cold water, and fanning the casualty to help it evaporate.
  - Placing in a cool shower if they are conscious enough to do so.
  - Spraying with cool water from a garden hose.
- If the casualty has a seizure, treat as you would for a febrile convulsion (*page 49*).



*Cool the casualty rapidly.*

## Taking a temperature

Modern, easy to use thermometers are now available, such as disposable strips that can be placed on the tongue or forehead. For these thermometers follow the manufacturers instructions. If you only have an 'old fashioned' mercury thermometer however, the following advice may help:

- Take care when handling the thermometer. The mercury centre is poisonous.
- Ensure that it has been properly cleaned.
- Hold the thermometer at the opposite end to the silver mercury bulb.
- Shake the thermometer until the mercury falls well below the 35°C mark.
- Place under the tongue of an adult (*who is fully conscious*), or the armpit of a child.
- Keep in place for 3 minutes.
- Read the temperature at the level to which the mercury has risen.

### 'Recreational' Drugs

*In recent years ambulance services have seen an increase in the use of 'recreational' drugs such as ecstasy (or 'e').*

*A casualty under the influence of such a drug may dance continually for long periods, which causes them to sweat excessively, and thus become hot and dehydrated.*

*The effects of dehydration, combined with the drug affecting 'normal' thought, can lead to heat exhaustion and heat stroke.*

## Diabetes

Diabetes is the name for a condition suffered by a person who does not produce enough of a hormone called insulin.

Insulin breaks down the sugar that we digest, so that it can be used by the cells of the body or stored for later use. In summary, insulin reduces the amount of sugar in the blood.


If diabetes goes untreated, the level of sugar in the blood will climb dangerously high over 1 to 2 days (*depending on the severity of the condition*).

There are 3 different types of diabetes, which are categorised by their method of treatment:

**Diet Controlled** This patient still produces some insulin naturally, so can control the condition by reducing the amount of sugar that they eat.

**Tablet Controlled** This patient still produces a small amount of insulin naturally, but needs to take tablets to help reduce the level of sugar in the blood, as well as diet control.

**Insulin Dependent** This patient produces little or no insulin, and has to inject themselves with insulin 2 or more times a day in order to keep sugar levels under control.



## High blood sugar – (hyperglycaemia)

Hyperglycaemia is the condition that occurs if diabetes has not been treated effectively with the methods mentioned above.

The sugar levels in the blood become higher and acids build up. The signs and symptoms in the table (*opposite*) are as a direct result of the body trying to excrete this acid build up.

## Low blood sugar – (hypoglycaemia)

Low blood sugar occurs mainly with diabetic patients who are insulin dependent, because the level of insulin in the body is now a 'fixed' amount because it is injected.

Because the patient has injected this 'fixed' amount of insulin, they have to balance it with the amount of food that they eat.

The blood sugar levels will fall low if:

- The patient does not eat enough food.
- The patient over exercises (*burning off sugar*).
- The patient injects too much insulin.

## Why is low blood sugar dangerous?

Unlike other cells in the body, the brain can only use glucose (*sugar*) as its source of energy. If the sugar in the blood becomes low therefore, the brain cells are literally starved.

The signs and symptoms of low blood sugar in the table (*opposite*) are as a result of the hungry brain cells becoming disordered, and the release of adrenaline that the disorder in the brain causes. (*see also 'the body's response to hypoxia'- page 14*).

## Possible signs and symptoms ?

|                           | High Blood Sugar<br>(hyperglycaemia)   | Low Blood Sugar<br>(hypoglycaemia)  |
|---------------------------|--|---|
| <b>Onset</b>              | Slow – 12 to 48 hours  | Fast – 2 minutes to 1 hour  |
| <b>Levels of Response</b> | Deteriorate slowly during the onset: <ul style="list-style-type: none"> <li>• Drowsy, lethargic behaviour</li> <li>• Unconsciousness if the condition is left untreated</li> </ul> | Deteriorate rapidly: <ul style="list-style-type: none"> <li>• Weakness, dizziness</li> <li>• Confusion, memory loss</li> <li>• Lack of coordination</li> <li>• Slurred speech</li> <li>• Bizarre, uncharacteristic, uncooperative, possibly violent behaviour</li> <li>• Unconsciousness within 1 hour</li> </ul> |
| <b>Skin</b>               | Dry and warm   | Pale, cold and sweaty   |
| <b>Breathing</b>          | Deep sighing breaths   | Normal, or shallow and rapid  |
| <b>Pulse</b>              | Rapid  | Rapid   |
| <b>Other Symptoms</b>     | Excessive urination<br>Excessive thirst<br>Hunger<br>Fruity odour on the breath  | Beware – the signs and symptoms can be confused for drunkenness   |

## Treatment of high blood sugar +

- Arrange for the patient to see a doctor as soon as possible.
- If the patient becomes unconscious, maintain Airway and Breathing, and dial 999 for an ambulance (see pages 6 to 8).

## Treatment of low blood sugar +

### For a conscious casualty:

- Sit the casualty down.
- Give the casualty a sugary drink (*isotonic sports drinks are best*), sugar lumps, glucose tablets, chocolate, or other sweet foods.
- If the casualty responds to treatment quickly, give them more food or drink.
- Stay with the casualty and let them rest until the level of response is 'fully alert' (see page 9).
- Tell the patient to see their doctor – even though they have fully recovered.
- If the patient does not respond to treatment within 10 minutes, or they are unmanageable, dial 999 for an ambulance.
- Consider if there is another cause for the patient's symptoms.

### For an unconscious casualty:

- Open the Airway and check for Breathing. Resuscitate as necessary (pages 6 to 8).
- Place the casualty in the recovery position if they are breathing effectively.
- Dial 999 for an ambulance.



Give the casualty a sugary drink – isotonic sports drinks are best.

**DO NOT** attempt to give the casualty anything to eat or drink if they become unconscious.

## Epilepsy

A person diagnosed with epilepsy has a tendency to have recurrent seizures (*fits*) that arise from a disturbance in the brain. This chapter does not only deal with patients who are diagnosed with epilepsy however, because one person in 20 will have a seizure at some point in their lives.

There are many causes of seizure (*including epilepsy*), such as hypoxia, stroke, head injury or even the body's temperature becoming too high.

Babies and young children commonly suffer seizures from becoming too hot due to illness and fever. This is covered in the topic 'febrile convulsions', opposite.

## Minor seizures

Minor epilepsy is also known as 'absence seizures' or 'petit mal' seizures. The patient may appear to suddenly start day dreaming (*even mid sentence*). This may last just a few seconds before recovery, and the patient might not even realise what has happened. Sometimes a minor seizure may be accompanied by unusual movements, such as twitching the face, jerking of an individual limb, or lip smacking. The patient may make a noise, such as letting out a cry.

## Treatment of minor seizures



- Remove any sources of danger, such as a knife or hot drink in their hands.
- Help the patient to sit down in a quiet place and reassure them.
- Stay with the patient until they are fully alert (*page 9*).
- If the patient is unaware of their condition, advise them to see a doctor.

## Major seizures

This type of seizure results from a major disturbance in the brain, which causes aggressive fitting, usually of the whole body.

Witnessing a major seizure can be frightening for the first aider, but calm, prompt action is essential for the patient.

## Possible signs and symptoms



A major seizure usually goes through a pattern:

- |                       |  |
|-----------------------|--|
| <b>Aura</b>           | If the patient has had seizures before, they may recognise that they are about to have one. The warning sign may be anything from a strange taste in the mouth, a smell, or a peculiar feeling. The aura may give the patient chance to seek help, or simply lie down before they fall.  |
| <b>'Tonic' Phase</b>  | Every muscle in the body suddenly becomes rigid. The patient may let out a cry and will fall to the floor. The back may arch and the lips may go blue ( <i>cyanosis</i> ). This phase typically lasts less than 20 seconds.  |
| <b>'Clonic' Phase</b> | The limbs of the body make sudden, violent jerking movements, the eyes may roll, the teeth may clench, saliva may drool from the mouth ( <i>sometimes blood-stained as a result of biting the tongue</i> ) and breathing could be loud like 'snoring'. The patient may lose control of the bladder or bowel.<br><br>This phase can last from 30 seconds to hours, although most seizures stop within a couple of minutes. Any seizure ( <i>or series of seizures</i> ) lasting more than 15 minutes is a dire medical emergency. |
| <b>Recovery Phase</b> | The body relaxes, though the patient is still unresponsive. Levels of response ( <i>page 9</i> ) will improve within a few minutes, but the patient may not be 'fully alert' for 20 minutes or so. They may be unaware of their actions and might want to sleep to recuperate.   |

## Treatment of major seizures (*fitting*)



### During the seizure:

- Help the patient to the floor to avoid injury if possible.
- Gently cushion the patient's head to help avoid injury. This can be done simply with your hands or a folded coat.
- Loosen any tight clothing around the neck to help the patient breathe.
- Move any objects from around the patient that may harm them and ask bystanders to move away.
- If you are concerned about the Airway, roll the casualty onto their side.
- Take note of the exact time the seizure started and its duration.
- Look for identification if you don't know the patient.



*Gently protect the head.*

### Dial 999 for an ambulance if:

- The seizure lasts more than 3 minutes.
- The patient's levels of response (*page 9*) don't improve after the seizure within 10 minutes.
- The patient has a second seizure.
- The patient is not diagnosed as epileptic or this is their first seizure.
- You are unsure.

**NEVER** place anything in the casualty's mouth (especially your fingers!).

**NEVER** try to hold the patient down or restrain them.

**NEVER** move the casualty (unless they are in danger).

### As soon as the seizure stops:

- Check Airway and Breathing. Resuscitate if necessary (*pages 6 to 8*).
- Place the patient in the recovery position (*page 12*).
- Keep the patient warm (unless temperature caused the seizure) and reassure them.
- Monitor Airway and Breathing.
- Move bystanders away before the casualty awakes and protect modesty.
- Check the levels of response regularly (*page 9*). Dial 999 if they don't improve within 10 minutes (or for any of the reasons mentioned above).

## Febrile convulsions

In young children and babies the area of the brain that regulates temperature (*the hypothalamus*) is not yet fully developed. This can lead to the core temperature of the body reaching dangerously high levels (*page 42*) and commonly a child in this situation may fit.

A febrile convulsion can be very frightening for the parents of the child. During the 'tonic' phase of the fit (*page 48*) the child may stop breathing, because the diaphragm goes rigid, and the lips and face may go blue (*cyanosis*). It goes without saying therefore, that calm reassurance will be necessary.

The child may have been unwell over the past day or so and will be hot to touch.

## Treatment of febrile convulsions



- Remove clothing and bedclothes. Provide fresh, cool air to cool the child down. Take care not to cool the child too much.
- Place the child on their side if possible to protect the Airway.
- Remove nearby objects and use padding to protect the child from injury whilst fitting. Pay particular attention to protecting the head.
- Dial 999 for an ambulance.
- If the child is still fitting – sponge them with tepid water to help the cooling process, but take care not to cool them too much.
- Constantly monitor Airway and Breathing until the ambulance arrives.



*If the child is still fitting – protect the child from injury and sponge with tepid water.*

**The digestive system**

Food enters the body through the mouth, where it is mechanically broken down by chewing, and the salivary glands secrete saliva, which helps break down starches (amongst other substances).

As we swallow, the epiglottis folds down to prevent food entering the airway, and the 'bolus' of food enters the oesophagus.

The bolus of food is pushed through the oesophagus (and the rest of the digestive system) by waves of muscle contractions.

Food enters the stomach, where acidic gastric juices are secreted to help break down the bolus of food to a soup like consistency.

The food then enters the duodenum, which is a duct into which enzymes from the pancreas, gall bladder and liver are secreted. These enzymes enable food to be broken down further, as it continues into the small intestine.

Although it is called the 'small' intestine, this duct is around 5 metres in length, and coils around in the centre of the abdominal cavity.

The small intestine completes the digestion process by absorbing nutrients from the food into the blood stream for use by the body.

Undigested food now passes into the large intestine (colon), where water is absorbed into the body, before being excreted from the anus.


Up to 1 minute

30 seconds

2 to 4 hours

2 to 6 hours

10 hours to several days





## Health and safety (first aid) regulations 1981

### Employer's responsibilities

Under Health and Safety law, an employer has a responsibility to ensure that first aid provision in the workplace is sufficient. This includes:

- Carrying out an assessment to decide where, how many and what type of First Aiders are needed, following guidance from the Health and Safety Executive.
- Providing training and refresher training for those First Aiders.
- Providing sufficient first aid kits and equipment for the workplace.
- Ensuring that all staff are aware of how and where to get first aid treatment.

This chapter gives some guidance on these responsibilities, although first aid training organisations are always willing to give advice.

### First aid kits

First Aid kits should be easily accessible, preferably placed near to hand washing facilities and clearly identified by a white cross on a green background. The container should protect the contents from dust and damp.

A first aid kit should be available at every work site. Larger sites may need more than one first aid kit. The following list of contents is given as guidance:

- 1 leaflet giving general guidance on first aid.
- 20 individually wrapped plasters of assorted size and appropriate to the type of work (e.g. blue detectable plasters should be provided for food handlers). Hypoallergenic plasters can be provided if necessary.
- 2 sterile eye pads.
- 4 triangular bandages, individually wrapped and preferably sterile.
- 6 safety pins.
- 6 medium wound dressings (approx. 12cm x 12cm), individually wrapped and sterile.
- 2 large wound dressings (approx. 18cm x 18cm), as above.
- 1 pair of disposable gloves.



The list is not mandatory, so equivalent items may be used. Additional items may be required, such as scissors, adhesive tape, disposable aprons and individually wrapped moist wipes. They may be stored in the first aid kit if they will fit, or kept close by for use.

Other items that may need to be considered are such things as blankets to protect casualties from the elements, or protective equipment such as breathing apparatus if a First Aider had to enter a dangerous atmosphere.

### Eye wash

If mains tap water is not readily available for eye irrigation, at least 1 litre of sterile water or 'saline' should be provided in sealed disposable container(s).

### Travelling first aid kits

First Aid kits for travelling workers should typically include:

- 1 leaflet giving general guidance on first aid.
- 6 individually wrapped sterile plasters.
- 1 large wound dressing (approx. 18cm x 18cm).
- 2 triangular bandages.
- 2 safety pins.
- Individually wrapped moist cleansing wipes.
- 1 pair of disposable gloves.

## First aid needs assessment

All employers must carry out a first aid needs assessment which should consider:

- the nature of the work and workplace hazards and risks.
- the size of the organisation.
- the nature of the workforce.
- the organisation's history of accidents and illness.
- the needs of travelling, remote and lone workers.
- work patterns such as shift work.
- the distribution of the workforce.
- the remoteness of the site from emergency medical services.
- employees working on shared or multi-occupied sites.
- annual leave and other absences of first aiders.
- first-aid provision for non-employees.

## Workplace hazards and risks

One of the more complicated areas of the first aid needs assessment is considering 'the nature of the work and workplace hazards and risks'.

An employer should consider the risks and identify what possible injuries could occur in order to ensure sufficient first aid provision is available.

The following table, compiled using information from the Health and Safety Executive, identifies some common workplace risks and the possible injuries that could occur:

| Risk                  | Possible Injuries Requiring First Aid  |
|-----------------------|--|
| Manual Handling       | Fractures, lacerations, sprains and strains.                                       |
| Slip and trip hazards | Fractures, sprains and strains, lacerations.                                       |
| Machinery             | Crush injuries, amputations, fractures, lacerations, eye injuries.                 |
| Work at height        | Head injury, loss of consciousness, spinal injury, fractures, sprains and strains. |
| Workplace transport   | Crush injuries, fractures, sprains and strains, spinal injuries.                   |
| Electricity           | Electric shock, burns.   |
| Chemicals             | Poisoning, loss of consciousness, burns, eye injuries.                             |

The employer should use these considerations to ensure that the correct level of first aider is trained to deal with the possible injuries or illness that could occur in the workplace (*see below*).

## First aiders

The selection of a first aider depends upon a number of factors. The person best suited to be a first aider will volunteer, and will have:

- good reliability, disposition and communication skills.
- an aptitude and ability to absorb new skills and knowledge.
- an ability to cope with stressful and physically demanding emergency procedures.
- normal duties in the workplace that can be left, to respond immediately and rapidly to an emergency.

From October 2009 there is a new training regime for first aiders. The Health and Safety Executive have introduced two levels of first aider;

- First Aider at Work; and
- Emergency First Aider at Work.

## Contents of HSE First Aid Courses:

| EFAW = Emergency First Aid at Work (1 day course)<br>FAW = First Aid at Work (3 day course)    | EFAW<br>1 day<br>(6 hours) | FAW<br>3 days<br>(18 hours) |
|--|----------------------------|-----------------------------|
| Acting safely, promptly and effectively in an emergency  | •                          | •                           |
| Cardio Pulmonary Resuscitation (CPR)   | •                          | •                           |
| Treating an unconscious casualty (including seizure)   | •                          | •                           |
| Wounds and bleeding  | •                          | •                           |
| Shock  | •                          | •                           |
| Minor injuries   | •                          | •                           |
| Choking  | •                          | •                           |
| Preventing cross infection, recording incidents and actions and the use of available equipment | •                          | •                           |
| Fractures  |                            | •                           |
| Sprains and strains  |                            | •                           |
| Spinal injuries  |                            | •                           |
| Chest injuries   |                            | •                           |
| Severe burns and scalds  |                            | •                           |
| Eye injuries   |                            | •                           |
| Poisoning  |                            | •                           |
| Anaphylaxis  |                            | •                           |
| Heart attack   |                            | •                           |
| Stroke   |                            | •                           |
| Epilepsy   |                            | •                           |
| Asthma   |                            | •                           |
| Diabetes   |                            | •                           |


These topics are not covered on the HSE Emergency First Aid at Work syllabus. The employer should ensure that the correct level of training is provided for first aiders to ensure that they can deal with the possible injuries or illness that could occur in the workplace.

**NOTE:** If the first aid needs assessment identifies the need for First Aid at Work (3 day) training, it is not acceptable to provide Emergency First Aiders.

## Annual refresher training

The HSE now recommend that all First Aiders and Emergency First Aiders attend annual refresher training due to the wealth of evidence on the severity of 'first aid skill fade'.

The flow chart, above right, shows the HSE's recommended sequence of training.



## Reporting of incidents at work

Any accident at work, no matter how small, must be recorded in an accident book (see overleaf). The incident may also need to be reported directly to the Health & Safety Executive under RIDDOR regulations:

### RIDDOR 1995 regulations

Reporting of Injuries, Diseases and Dangerous Occurrences Regulations 1995.

These regulations state that it is the responsibility of the *Employer or person in control of the premises* to report the following occurrences directly to the Health and Safety Executive:

- Deaths (report immediately).
- Major injuries (report immediately).
- Dangerous occurrences (report immediately).
- Incidents resulting in a person being off work (or unable to do full duties) for more than 3 days (report within 10 days).
- Diseases (report as soon as possible).

For more information you can visit the RIDDOR website at: [www.riddor.gov.uk](http://www.riddor.gov.uk)



## Accident book

Any accident at work, no matter how small, must be recorded in an accident book. The accident book may be filled in by any person on behalf of the casualty (*or indeed by the casualty themselves*).

The information recorded can help the employer identify accident trends and possible areas of improvement in the control of health and safety risks. It can be used for future first aid needs assessments and may be helpful for insurance investigative purposes.

Filling in the accident book is often done by the First Aider, so the following notes are given for your advice:

- An accident book is a legal document.
- Anything that has been written down at the time of an incident is usually considered to be 'stronger evidence' in court than something recalled from memory.
- Complete the report all at the same time, using the same pen (*not pencil*).
- To comply with the Data Protection Act, personal details entered in accident books must be kept confidential, so the book should be designed so that individual record sheets can be removed and stored securely.
- A member of staff should be nominated to be responsible for the safekeeping of completed accident records (*e.g. in a lockable cabinet*). Hand the completed accident record to that person.
- The person who had the accident may wish to take a photocopy of the report. If this is the case, they can do this before it is handed in. They should keep a record of the accident report number.

A typical accident record form.

### You should include in the report:

- The name, address and occupation of the person who had the accident.
- The name, address, occupation and signature of the person who is completing the report.
- The date, time and location of the accident.
- A description of how the accident happened, giving the cause if you can.
- Details of the injury suffered.

## First aid patient report form

It is useful for a first aider to complete a patient report form for every patient treated. Please note that this does not replace the accident book, which would still have to be completed for an accident at work.

A copy of a patient report form is opposite. You can make copies of this for your own use.

The patient report form is designed so the first aider can keep a record of the exact treatment provided. It is particularly useful if a patient refuses treatment against the advice of the first aider.

- If a patient refuses treatment, make sure they are capable of making that decision (*e.g. a fully conscious adult*). Seek medical advice if they are not.
- Follow the advice given for completing the accident book (*above*) when completing the form.
- A copy of the form can be given to ambulance or hospital staff, as it will contain valuable information about the incident and treatment of the patient. Ask the nurse to take a copy, so you can keep the original.
- To comply with the Data Protection Act, personal details on the report form must be kept confidential, so the report should be stored securely (*e.g. in a lockable cabinet*).

## AVPU score

A simple way to record the conscious level of a patient is to use the 'AVPU' scale. A detailed explanation of the scale is given on page 9.

The scale is listed on the patient report form so you don't have to remember it. There is a score provided next to each level of consciousness. Write the score in the observations chart each time you measure it.

# First aid patient report form

Date \_\_\_\_\_ Time \_\_\_\_\_ First Aider \_\_\_\_\_

Patient Name \_\_\_\_\_ Sex \_\_\_\_\_ Date of Birth \_\_\_\_\_ Age \_\_\_\_\_

Patient's Address / Job details \_\_\_\_\_


Location of Incident \_\_\_\_\_ Time of incident \_\_\_\_\_

**Patient Observations:** (record at least every 10 minutes)

| Time |  |  |  | Breathing Rate |  | Pulse Rate |  | AVPU SCORE |
|------|--|--|--|----------------|--|------------|--|------------|
|      |  |  |  |                |  |            |  |            |
|      |  |  |  |                |  |            |  |            |
|      |  |  |  |                |  |            |  |            |
|      |  |  |  |                |  |            |  |            |
|      |  |  |  |                |  |            |  |            |
|      |  |  |  |                |  |            |  |            |
|      |  |  |  |                |  |            |  |            |

| AVPU SCORE:          |  | SCORE: |
|----------------------|--|--------|
| <b>A</b> lert        | Fully Alert (usually knows the month).   | 6      |
| <b>V</b> oice        | Confused.                                | 5      |
|                      | Inappropriate Words.                     | 4      |
|                      | Utters Sounds.                           | 3      |
| <b>P</b> ain         | Localises Pain.                          | 2      |
|                      | Responds to (but does not localise) Pain | 1      |
| <b>U</b> nresponsive | Unresponsive to speech and pain stimuli. | 0      |

**A.M.P.L.E.**

|                            |   |
|----------------------------|---|
| Allergies                  |   |
| Medication                 |   |
| Past Medical History       |   |
| Last Eaten                 |   |
| Events Leading to Incident |   |
| Treatment / Comments       |    |
| What happened afterwards?  | <p><b>The patient went:</b>    <input type="checkbox"/> home    <input type="checkbox"/> to hospital    <input type="checkbox"/> to GP    <input type="checkbox"/> in ambulance    <input type="checkbox"/> back to work</p> <p>other / details _____</p> |

Patient's Signature: \_\_\_\_\_

Date: \_\_\_\_\_

First Aider's Signature: \_\_\_\_\_

Date: \_\_\_\_\_



Gently tap the shoulders and shout.



Open the airway.



Check for normal breathing.



Rescue breaths.



Use 1 or 2 hands to depress the chest by one third of its depth.

**NOTE:** The minor modifications to the adult sequence of resuscitation (page 8) that can make it even more suitable for children have been included below.

**REMEMBER:** If you are unsure, it is better to perform the adult sequence of resuscitation on a child (who is unresponsive and not breathing) than to do nothing at all.

## Child resuscitation

### **D** Danger

- Check that it is safe for you to help. Do not put yourself at risk.

### **R** Response

- Gently tap the shoulders and shout 'are you alright?'
- If there is no response, shout for help, but don't leave the child just yet.

### **A** Airway

Carefully open the airway by using 'head tilt' and 'chin lift':

- Place your hand on the forehead and gently tilt the head back.
- With your fingertips under the point of the chin, lift the chin to open the airway.

### **B** Breathing

Keeping the airway open, look, listen and feel to see if the breathing is normal. Take no more than 10 seconds to do this.

- If the child is breathing **normally**, carry out a secondary survey and place them in the recovery position (page 11).

#### If the child is not breathing normally:

- Ask someone to dial 999 for an ambulance immediately, but if you are on your own, and you have to leave the child to dial 999, carry out resuscitation for about 1 minute first:
- Keep the airway open by tilting the head and lifting the chin.
- Nip the nose and seal your mouth around the child's mouth.
- Give 5 initial rescue breaths (*blow in just enough air to make the child's chest visibly rise*).

#### Combine rescue breaths with chest compressions:

- Use 1 or 2 hands as required to depress the chest a third of its depth.
- Give 30 chest compressions at a rate of 100 per minute.
- Open the airway again by tilting the head and lifting the chin, then give 2 more rescue breaths.
- Continue repeating cycles of 30 compressions to 2 rescue breaths.
- Only stop to recheck the child if they start breathing normally – otherwise do not interrupt resuscitation.

#### If your rescue breaths don't make the chest rise effectively:

Give another 30 chest compressions, then before your next attempt:

- Check inside the mouth and remove any visible obstruction (*but don't reach blindly into the back of the throat*).
- Recheck there is adequate head tilt and chin lift.
- Do not attempt more than 2 breaths each time before returning to chest compressions.

**NOTE:** If there is more than one rescuer, change over every two minutes to prevent fatigue. Ensure the minimum of delay as you change over.

**NOTE:** The minor modifications to the adult sequence of resuscitation (page 8) that can make it even more suitable for children and babies have been included below.

**REMEMBER:** If you are unsure, it is better to perform the adult sequence of resuscitation on a baby (who is unresponsive and not breathing) than to do nothing at all.

## Baby resuscitation



### D Danger

- Check that it is safe for you to help. Do not put yourself at risk.

### R Response

- Gently tap the shoulders and shout to try and wake the baby.
- If there is no response, shout for help, but don't leave the baby just yet.

### A Airway

Carefully open the airway by using 'head tilt' and 'chin lift':

- Place your hand on the forehead and gently tilt the head back. Do not over-extend the neck.
- With your fingertips under the point of the chin, gently lift the chin to open the airway.

### B Breathing

Keeping the airway open, look, listen and feel to see if the breathing is normal. Take no more than 10 seconds to do this.

- If the baby is breathing normally, consider injuries and place them in the recovery position (page 11).

#### If the baby is not breathing normally:

- Ask someone to dial 999 for an ambulance immediately, but if you are on your own, and you have to leave the baby to dial 999, carry out resuscitation for about 1 minute first:
- Keep the airway open by tilting the head and lifting the chin (*do not over-extend the neck*).
- Seal your mouth around the baby's mouth and nose.
- Give 5 initial rescue breaths (*blow in just enough air to make the chest visibly rise*). Take care not to over inflate the lungs.

#### Combine rescue breaths with chest compressions:

- Use 2 fingers to depress the chest a third of its depth.
- Give 30 chest compressions at a rate of 100 per minute.
- Open the airway again by tilting the head and lifting the chin, then give 2 more rescue breaths.
- Continue repeating cycles of 30 compressions to 2 rescue breaths.
- Only stop to recheck the child if they start breathing normally – otherwise don't interrupt resuscitation.

#### If your rescue breaths don't make the chest rise effectively:

Give another 30 chest compressions, then before your next attempt:

- Check inside the mouth and remove any visible obstruction (*but don't reach blindly into the back of the throat*).
- Recheck there is adequate head tilt and chin lift.
- Do not attempt more than 2 breaths each time before returning to chest compressions.



Open the airway.



Check for normal breathing.



Rescue breaths.



Use 2 fingers to depress the chest by one third of its depth.



Some examples of AEDs.

## Automated External Defibrillation



The most common cause for a heart to stop (*cardiac arrest*) is a 'heart attack' (page 24). It is worth noting that a heart attack does not always cause a cardiac arrest. The majority of people who suffer a heart attack stay conscious and survive.

If a heart attack (*or other cause*) results in a cardiac arrest, it is usually because it has interrupted the heart's electrical impulses. When this happens the heart quivers chaotically instead of beating in a co-ordinated rhythm. This is called 'Ventricular Fibrillation' (VF).

The definitive treatment of Ventricular Fibrillation is to deliver a controlled electric shock through the heart, to stop the 'quivering' and enable it to beat normally again. This is called 'defibrillation'.

An Automated External Defibrillator (AED) is a safe, reliable, computerised device that can analyse heart rhythms and enable a non-medically qualified rescuer to safely deliver the life saving shock with a small amount of training.

The use of an AED can dramatically increase the chances of survival if a patient's heart stops beating, but it must be used promptly – for every one minute delay in delivering the shock, the patient's chance of survival reduces by up to 10%.



## Resuscitation with an AED



### D Danger

- Check that it is safe for you to help the casualty. Do not put yourself at risk.
- Consider the safety implications of using an AED in this situation (page 60).

### R Response

- Gently shake the shoulders and ask loudly 'Are you alright?'

If there is no response:

- Shout for help immediately.
- If possible, ask one helper to dial 999 and another to get the AED, but do not leave the casualty yourself just yet.

### A Airway

- Carefully open the airway by using 'head tilt' and 'chin lift'.

### B Breathing

Keeping the airway open, look, listen and feel to see if the breathing is normal. Take no more than 10 seconds to do this.

- If the casualty is breathing normally, consider possible injuries and carefully place them in the recovery position (page 11).

#### If the casualty is not breathing normally:

- If you are on your own, dial 999 for an ambulance and get the AED – you may need to leave the casualty to do this.
- If you have help – start CPR immediately whilst your helper(s) get the AED and dial 999. Continue CPR until the AED arrives (see pages 6 and 7).



Open the airway.



Look, listen and feel for normal breathing.




## When the AED arrives:

- If you have a helper, ask them to continue CPR whilst you get the AED ready.  
**NOTE:** *If the helper is untrained it may be easiest for them to give chest compressions only – see page 8.*

### Switch on the AED immediately and follow the voice prompts:

- Attach the leads to the AED if necessary and attach the pads to the victim's bare chest (*do this whilst your helper performs CPR if possible*).
  - You may need to towel dry or shave the chest so the pads adhere properly. Only shave excessive hair and don't delay defibrillation if a razor is not immediately available.
  - Peel the backing from one pad at a time and place firmly in position, following the instructions on the pads.
  - Place one pad below the victim's right collar bone.
  - Place the other pad on the victim's left side, over the lower ribs. Place this pad vertically if possible (*see below*).
- DO NOT** remove the pads if you have placed them the wrong way around – the AED will still work.
- Whilst the AED analyses the rhythm – stop CPR and ensure that no one touches the casualty.



## If a shock is advised:

- Ensure that nobody touches the casualty (*check from top to toe and shout 'stand clear!'*)
- Push the shock button as directed (*fully-automatic AEDs will deliver the shock automatically*).
- Continue as directed by the voice / visual prompts.

## If a shock is NOT advised:

- Immediately resume CPR using a ratio of 30 chest compressions to 2 rescue breaths.
- Continue as directed by the voice / visual prompts.

## Placement of the pads



### Wet chest

If the patient's chest is wet (*perfuse sweating for example*), it must be dried before applying the pads so they adhere to the chest properly. Also dry the chest between the pads so electricity does not 'arc' across the wet chest.



### Excessive chest hair

Chest hair will prevent the pads adhering to the skin and will interfere with electrical contact. Only shave the chest if the hair is excessive and even then spend as little time as possible on this. Do not delay defibrillation if a razor is not immediately available.



### Pad positioning

Recent studies have found that the position of the pad on the lower left chest can affect the effectiveness of the shock. Ensure the pad is placed around the side of the chest (*not on the front*) and place it vertically (*see picture*). This ensures the maximum electricity flows through the heart rather than across the chest surface. Until manufacturers update them, some AED pads will have diagrams showing horizontal placement – ignore this and place the pad vertically.



*Pads too close together: Electricity flows across the chest.*



*Correct pad positions: Electricity flows through the heart.*



*Lower left pad positioned vertically.*

## AED safety considerations



### Electrical shock

Recent tests have shown that if the patient's chest is dry and the pads are stuck to the chest correctly, the risk of electrical shock is very low, because the electricity only wants to travel from one pad to the other, not to 'earth' like mains electricity. To be extra safe however, briefly check that nobody is touching the patient before delivering a shock.

**DO NOT** delay defibrillation because the patient is on a wet or metal surface – providing the chest is dry it is safe to deliver the shock.

### Medication patches

Some patients wear a patch to deliver medication (such as a nicotine patch). Some heart patients wear a 'glyceryl tri-nitrate' (GTN) patch. This type of patch can explode if electricity is passed through it, so remove any visible medication patches as a precaution before delivering the shock.




### Highly flammable atmosphere

There is a danger of the AED creating a spark when the shock is delivered, so it should not be used in a highly flammable atmosphere (in the presence of petrol fumes for example).

### Jewellery

Take care not to place the pads over jewellery such as a necklace. This would conduct the electricity and burn the patient. There is no need to remove pierced jewellery, but try to avoid placing the pads over it.



### Implanted devices

Certain heart patients may have a pacemaker or defibrillator implanted. You can often see or feel them under the skin when the chest is exposed and there may be a scar. They are usually implanted just below the left collar bone, which is not in the way of the AED pads, but if a device has been implanted elsewhere, try to avoid placing the pad directly over it.



### Inappropriate shock

AEDs are proven to analyse heart rhythms extremely accurately, however the patient needs to be motionless whilst the AED does this. Do not use an AED on a patient who is fitting (violent jerking movements) and ensure vehicle engines or vibrating machinery are switched off whenever possible.



## AED use on children



Normal 'adult' AED pads are suitable for a child older than 8 years. Smaller pads that reduce the current delivered are available for children aged 1 to 8. These should be used for that age range where available. Some AEDs have a 'paediatric' setting.

If the child is older than 1 year and you only have adult pads, use the AED as it is. The use of adult pads on a child under 1 year is not recommended.

Most paediatric pads are designed to be placed with one pad in the centre of the child's chest and the other pad in the centre of the back. The pads will have a diagram showing the correct positions.



Some paediatric pads are designed to be placed in the same location as adult pads. Always follow the diagrams on the pads to ensure correct placement.

|                                   |   |                                 |   |
|-----------------------------------|---|---------------------------------|---|
| <b>Abdomen</b>                    | The area between the lowest ribs and the pelvis.  | <b>Hepatic</b>                  | Relating to the liver.  |
| <b>Acute</b>                      | Sudden onset.   | <b>Hyper...</b>                 | High.   |
| <b>Adrenaline</b>                 | Hormone secreted by the body in times of shock ( <i>see page 14</i> ).                    | <b>Hypo...</b>                  | Low.  |
| <b>...aem...</b>                  | Referring to the blood.   | <b>Hypothalamus</b>             | Area of the brain that controls body temperature.   |
| <b>Airway</b>                     | The passage from the mouth and nose to the lungs.   | <b>Hypovolaemic</b>             | Low volume of blood, a type of shock.   |
| <b>Alveoli</b>                    | Minute air sacks in the lungs, through which the exchange of gases takes place.           | <b>Hypoxia</b>                  | Low levels of oxygen in the blood.  |
| <b>Asphyxia</b>                   | Deficiency of oxygen caused by an interruption in the passage of air to the lungs.        | <b>Inferior</b>                 | Below.  |
| <b>Atrium</b>                     | Top, 'collecting' chamber of the heart ( <i>of which there are two</i> ).                 | <b>Insulin</b>                  | Hormone secreted by the pancreas that enables the usage and storage of sugar.   |
| <b>Baby</b>                       | Person under 1 year old.  | <b>Jaw Thrust</b>               | Manoeuvre to open the airway without moving the head, by thrusting the jaw forwards.  |
| <b>Breathing</b>                  | Inspiration and expiration of air into and out of the lungs.                              | <b>Mesenteric</b>               | Relating to an area of the intestines.  |
| <b>Bronchioles</b>                | Small air passages in the lungs, leading to the alveoli.                                  | <b>Nausea</b>                   | Feeling sick.   |
| <b>Cardiac / cardiogenic</b>      | Concerned with the heart.   | <b>Neurogenic</b>               | Concerned with the brain and nervous system.  |
| <b>Cell</b>                       | Smallest structural living unit of an organism.   | <b>Perfusion</b>                | Supply of oxygen and nutrients, and the removal of waste gases and products.  |
| <b>Cerebro-spinal fluid (CSF)</b> | Fluid that surrounds the brain and spinal cord, to cushion it and provide nutrients etc.  | <b>Pleura</b>                   | A two layered membrane surrounding the lungs, between which is a 'serous' fluid.  |
| <b>Cerebrum</b>                   | The largest part of the brain.  | <b>Pneumothorax</b>             | Air entry into the pleural cavity of the lung.  |
| <b>Cervical</b>                   | Concerned with the neck.  | <b>Pulmonary</b>                | Concerned with the lungs.   |
| <b>Child</b>                      | Person between 1 year old and puberty.  | <b>Regurgitation</b>            | Vomiting, being sick.   |
| <b>Chronic</b>                    | Long term.  | <b>Rescue Breath</b>            | Blowing air into a patient's lungs, sufficient to make the chest rise.  |
| <b>Circulation</b>                | The movement of blood around the body.  | <b>Respiration</b>              | Breathing.  |
| <b>Compression</b>                | Bleeding or swelling in the cranial cavity, exerting pressure on the brain.               | <b>Seizure</b>                  | Fit or Convulsion.  |
| <b>Concussion</b>                 | Shaking of the brain, causing temporary loss of consciousness or function.                | <b>Shock</b>                    | Inadequate supply of oxygen to the tissues as a result of a fall in blood pressure or volume.                                     |
| <b>Consciousness</b>              | Alertness, 'normal' activity of the brain.  | <b>Spinal cord</b>              | Group of nerves which emanate from the brain and pass down the spinal column.   |
| <b>Constrict</b>                  | To close down, become narrower.   | <b>Spine</b>                    | The column of vertebrae which form the back.  |
| <b>Convulsion</b>                 | Fit or Seizure.   | <b>Stroke</b>                   | Bleed or blockage of a blood vessel within the brain ( <i>see page 13</i> ).  |
| <b>CPR</b>                        | Cardio Pulmonary Resuscitation. Manually squeezing the heart and breathing for a patient. | <b>Superior</b>                 | Above.  |
| <b>Cranium</b>                    | The cavity in the skull in which the brain lies.  | <b>Symptoms</b>                 | The feelings of a patient e.g. "I feel sick."   |
| <b>Cyanosis</b>                   | Blue grey tinges to the skin, especially the lips, due to lack of oxygen.                 | <b>Syncope</b>                  | Faint.  |
| <b>Defibrillation</b>             | The delivery of a large electric shock to the chest in an attempt to re-start the heart.  | <b>Tension pneumothorax</b>     | Air entry into the pleural cavity of the lung that has become pressurised, impairing the function of the good lung and the heart. |
| <b>Dilate</b>                     | Become wider, open up.  | <b>Thoracic</b>                 | The area within the rib cage containing the lungs.  |
| <b>Enzyme</b>                     | Substance that enables a biological reaction to happen.                                   | <b>Tourniquet</b>               | A tight band placed around a limb which was used to stop blood flow. No longer used in first aid.                                 |
| <b>Epistaxis</b>                  | Nose bleed.   | <b>Ventricle</b>                | Lower, larger 'pumping' chamber of the heart ( <i>of which there are two</i> ).   |
| <b>Face shield</b>                | Protective mask with a one-way valve for performing mouth-to-mouth rescue breaths.        | <b>Ventricular Fibrillation</b> | Quivering, vibrating movement of the ventricles of the heart, producing no effective pumping action.                              |
| <b>Febrile</b>                    | Relating to fever or high body temperature.   |                                 |   |
| <b>Haemothorax</b>                | Bleeding into the pleural cavity of the lungs.  |                                 |   |

# First aid quiz

The following questions are provided so you can test your knowledge of first aid. Discuss with your first aid instructor which questions are best to revise. Write your answers on a separate sheet of paper.

Answer the questions as best as you can from memory, then have a look in the book to mark yourself (or improve your answers). The page numbers in *red italics* indicate where to look for the correct answer.

1. What is the order of priorities when dealing with a patient? *Page 4*
2. What ratio of chest compressions to rescue breaths would you do when performing CPR? *Page 7*
3. What modifications can you make to the adult sequence of CPR to make it even more suitable for a child? *Page 8*
4. What are the 2 main dangers facing someone who is unconscious and on their back. *Page 11*
5. How would you treat a patient who is unconscious because of head injury? *Page 13*
6. What are the signs and symptoms of hypoxia? *Page 14*
7. Try to remember as many of the causes of hypoxia as you can. *Page 14*
8. Someone starts to choke on some food. What should you do? *Pages 16 or 17*
9. Someone is suffering from anaphylaxis and they are struggling to breathe. How will they look? *Page 18*
10. How would you treat a patient having an asthma attack? *Page 19*
11. What are the average pulse rates of: a) an adult. b) a child. c) a baby. *Page 23*
12. What are the signs and symptoms of a heart attack? *Page 24*
13. How would you treat a patient suffering from heart attack? *Page 25*
14. Someone has cut their arm on some sharp metal. The metal isn't stuck in the wound but it is bleeding badly. What should you do? *Page 30*
15. Someone in the kitchen has slipped with a sharp knife and amputated their finger. Describe your actions. *Pages 30 and 32*
16. Someone has scalded the whole of their hand on some hot water. What should you do? What percentage of burn is that? *Pages 34 and 35*
17. Can you remember some signs and symptoms of a fracture? *Page 38*
18. A patient has fractured their wrist. How would you treat the patient? *Page 38*
19. A patient has fallen from a horse and they are unconscious. You check breathing and they are breathing normally. What should you do now? *Page 40*
20. Someone has been working outside in very cold rainy weather all day. You suspect they are hypothermic. How should you treat them? *Page 43*
21. A diabetic patient is suddenly acting strangely and not making sense. What is probably wrong? What should you do? *Pages 46 and 47*
22. How would you treat a patient having an epileptic seizure? What should you do after the seizure stops? *Page 49*

# First Aid Made Easy

- Updated with the latest resuscitation guidelines.
- Approved for use with HSE first aid at work courses.
- A logical and easy to understand layout.
- Simple, accurate information with lots of pictures and diagrams.
- Includes 'test yourself' questions.
- Written by an experienced Paramedic and First Aid Instructor.

RRP £6.99



**Mixed Sources**

Product group from well-managed  
forests and other controlled sources  
www.fsc.org Certno. SGS-COC-006290  
© 1996 Forest Stewardship Council



9 780955229404

ISBN-13: 978-0-9552294-0-4

ISBN-10: 0-9552294-0-5